



ADVIES VAN DE HOGE GEZONDHEIDSRAAD nr. 9216

Aanbevelingen over het gebruik van kunstmatige bronnen van UV-straling in België

In this scientific advisory report on public health policy, the Superior Health Council of Belgium provides recommendations for the Belgian population on the exposure to artificial UV radiation (sunbeds)

Versie gevalideerd op het College van
Juni - 2017¹

I INLEIDING EN VRAAGSTELLING

De Hoge Gezondheidsraad (HGR) heeft in 2000 een advies uitgebracht over zonnebanken (HGR 5783, 2000).

De gegevens in verband met deze problematiek zijn op dit ogenblik nog steeds geldig, maar zijn weliswaar niet meer up-to-date. Recente publicaties belichten de verschillende problemen, en vooral de aanzienlijke toename in het aantal gevallen van kanker die in verband worden gebracht met blootstelling aan ultraviolette straling (UV-straling), ook aan de UV-straling van zonnebanken.

De algemene bevolking en vooral jongeren die zich om cosmetische redenen in aanzienlijke mate aan UV-straling blootstellen, moeten over deze risico's worden ingelicht.

Bovendien moet de overheid over actuele informatie beschikken over de huidige situatie en over maatregelen die kunnen worden genomen om de gezondheidsrisico's van UV-straling in het algemeen en UV-straling geproduceerd door zonnebanken in het bijzonder, verder te doen dalen.

Om deze redenen heeft de HGR beslist om zijn vorige advies in het licht van de meest recente wetenschappelijke gegevens en de huidige (internationale) aanbevelingen te herzien.

¹ De Raad behoudt zich het recht voor om in dit document op elk moment kleine typografische verbeteringen aan te brengen. Verbeteringen die de betekenis wijzigen, worden echter automatisch in een erratum opgenomen. In dergelijk geval wordt een nieuwe versie van het advies uitgebracht.

II AANBEVELINGEN

De HGR heeft recente gegevens onderzocht om zijn advies over zonnebanken te updaten (HGR 5783, 2000).

Er werd een *ad hoc*-werkgroep opgericht met deskundigen in de volgende disciplines: risicoanalyse, dermatologie, psychologie, oogheelkunde en biofysica. Deze werkgroep heeft de literatuur over de gezondheidseffecten van UV-straling beoordeeld, met speciale aandacht voor zonnebanken voor cosmetische doeleinden.

Het is al jaren bewezen dat blootstelling aan UV-straling voor de aanmaak van vitamine D zorgt. Om het nodige gehalte aan vitamine D op te wekken is een beperkte blootstelling aan UV-straling echter voldoende en indien nodig bestaan er ook alternatieve bronnen voor vitamine D (orale vitamine D-supplementen).

Maar in de afgelopen decennia zijn er ook steeds meer aanwijzingen gevonden dat UV-straling, en ook de UV-straling van zonnebanken, zowel als initiator alsook als promotor van huidkanker optreedt. Meer concreet zijn er sterke bewijzen dat blootstelling aan UV-straling, ook de UV-straling van zonnebanken, op alle leeftijden het risico op het ontstaan van een cutaan melanoom en squameus celcarcinoom verhoogt en dat het risico op deze vormen van kanker (vooral melanoom) nog hoger is in geval van blootstelling tijdens de kindertijd en adolescentie. Uit een matig aantal gegevens blijkt bovendien ook dat blootstelling aan UV-straling, inclusief de UV-straling van zonnebanken, ook het risico op basaal celcarcinoom en oculair melanoom verhoogt.

Op basis van de solide bewijzen dat UV-straling, met inbegrip van de UV-straling van zonnebanken, huidkanker en andere huid- en oogziekten veroorzaakt, en rekening houdend met het feit dat

- er voor het opwekken van huidkanker geen drempelniveau bestaat voor UV-straling en dus ook geen veilige blootstellingsgrens;
- de risico's van blootstelling aan UV-straling zwaarder wegen dan de voordelen, zoals de aanmaak van vitamine D;
- een minimale blootstelling aan natuurlijke UV-straling doorgaans voldoende is om de nodige hoeveelheden vitamine D aan te maken;
- het gebruik van zonnebanken geen goede oplossing is om een vitamine D-tekort te verhelpen;
- zonnebanken permanent beschikbaar zijn, wat ertoe aanzet om deze herhaaldelijk en overmatig te gebruiken.

De blootstelling aan natuurlijke UV-straling kan aan de hand van beschermende maatregelen worden beperkt, maar niet volledig worden uitgesloten.

De HGR adviseert om zonnebanken en alle kunstmatige bronnen van UV-straling die beschikbaar zijn voor het publiek te verbieden, met als doel het risico op huidkanker te verminderen.

Tijdens de overgangperiode tot het verbod op zonnebanken effectief is, en ook achteraf, beveelt de HGR aan om de algemene bevolking op een objectieve en transparante manier in te lichten over de risico's van UV-straling in het algemeen (en ook over de risico's van de UV-straling die door zonnebanken wordt geproduceerd), en over de redenen achter het verbod op zonnebanken.

De bedoeling is om zo een verschuiving naar een verhoogde blootstelling aan natuurlijke UV-straling te vermijden. Bovendien is het ook belangrijk om in deze voorlichtingscampagne objectieve informatie op te nemen over de positieve effecten van incidentele blootstelling aan de zon, bv. de aanmaak van vitamine D.

Keywords and MeSH descriptor terms²

MeSH terms*	Keywords	Sleutelwoorden	Mots clés	Schlüsselwörter
Sunbed	Sunbed	Zonnebank	Banc solaire	
Ultraviolet radiation	Ultraviolet radiation	Ultraviolette straling	Rayonnement ultraviolet	
Skin Cancer	Skin cancer	Huidkanker	Cancer de la peau	

MeSH (Medical Subject Headings) is the NLM (National Library of Medicine) controlled vocabulary thesaurus used for indexing articles for PubMed <http://www.ncbi.nlm.nih.gov/mesh>.

² The Council wishes to clarify that the MeSH terms and keywords are used for referencing purposes as well as to provide an easy definition of the scope of the advisory report. For more information, see the section entitled "methodology".

I	INTRODUCTION AND ISSUE	1
II	RECOMMENDATIONS	2
III	METHODOLOGY	6
IV	ELABORATION AND ARGUMENTATION	7
1	Sources of UV radiation	7
1.1	Natural source (sunlight)	7
1.2	Artificial sources	8
1.2.1	Sunbeds and other tanning equipment	8
	Technical aspects	9
	Legal aspects	10
	Practice of sunbeds in Belgium	11
	Prevalence of sunbed use in Belgium	12
	Characteristics and behaviour of sunbed users in Belgium	13
	Knowledge and misconceptions	14
1.2.2	Other sources of UV radiation	15
2	Effects and health risks of UV	16
2.1	Intentional effects	16
2.1.1	Photo-protection effects	16
2.1.2	Sociological and psychological effects	18
	Sociological aspects	18
	Psychological aspects	19
2.1.3	Vitamin D	21
	Synthesis	21
	UV-lamps and endogenous vitamin D synthesis	21
	Health effects of vitamin D	23
	Vitamin D supplements	24
2.1.4	Other uses	21
2.2	Non-intentional effects	25
2.2.1	Carcinogenicity	25
	Genotoxicity	25
	Mutagenesis and Carcinogenesis	26
	Epidemiological studies (sunbed-related)	27
	Health risks	28
	Economic burden of skin cancer in Belgium	29
2.2.2	Ageing (heliodermy)	30
2.2.3	Other issues	31
	Ocular problems linked to UV radiation	31
	Photoallergic reactions	31
	Photoinduced disorders	31
	Photosensitisation reactions	32
V	CONCLUSION AND RECOMMENDATIONS	33
1	Conclusions	34
2	Recommendations	34
VI	REFERENCES	38
VII	COMPOSITION OF THE WORKING GROUP	45
VIII	APPENDIXES	47
1	Appendix 1: Definition and interpretation of the UV index	47
2	Appendix 2: Results of Euromelanoma survey (31 countries)	49

2.1	EU population	49
2.1.1	sunbed use	49
2.1.2	sunbed use (females < 35y)	50
2.2	Belgian population	51
2.2.1	Sunbed use (2015)	51
2.2.2	Number of years of use (2015)	51
2.2.3	Evolution of sunbed use (2009 – 2015)	51
2.2.4	Evolution of number of years of use	52

III METHODOLOGY

After determining the scope of the project, the Board and the Chair of the permanent working group “Cosmetology and cosmetic devices, including cosmetic surgery” identified the necessary fields of expertise. An *ad hoc* working group was then set up which included experts in risk analysis, dermatology, psychology, ophthalmology and biophysics. The experts of this working group provided a general and an *ad hoc* declaration of interests and the Committee on Deontology assessed the potential risk of conflicts of interest.

This advisory report is based on a review of the scientific literature published in both scientific journals and reports from national and international organisations competent in this field (peer-reviewed), as well as on the opinion of the experts.

Once the advisory report was endorsed by the *ad hoc* working group and by the standing working groups “Cosmetology and cosmetic devices, including cosmetic surgery” and “Non-ionising radiation” it was ultimately validated by the Board.

IV ELABORATION AND ARGUMENTATION

List of abbreviations used

ACGIH	American Conference of Governmental and Industrial Hygienists
FPS	Federal Public Service
GPSD	General Product Safety Directive
ICNIRP	International Commission on Non-Ionizing Radiation Protection
MC1R	melanocortin receptor 1
MSH	melanocyte-stimulating hormone
NMSC	non-melanoma skin cancer
POMC	pro-opiomelanocortin
SED	Standard Erythema Dose
SHC	Superior Health Council
SPF	sun protection factor
UV	ultraviolet
WMO	World Meteorological Organisation
WHO	World Health Organisation

1 Sources of UV radiation

Ultraviolet (UV)-radiation is a form of electromagnetic radiation, with wavelengths shorter than light (visible electromagnetic radiation), but longer than X-rays. UV radiation is usually classified as UV-A, UV-B and UV-C radiation (Table 1) ([Commission Internationale de l'Éclairage, 2009](#)).

Table 1: Classification of ultraviolet radiation ([Commission Internationale de l'Éclairage 2009](#)).³

Name	Wavelength range (nm)
UV-A	400-315
UV-B	315-280
UV-C	280-200

1.1 Natural source (sunlight)

The sun emits radiation across a very wide spectrum and is the only natural source of UV radiation. The Earth's atmosphere filters out a large part of the shortwave radiation. Therefore, almost no radiation with wavelengths shorter than 280 nm (UV-C) reaches the surface of the Earth. The absorption of radiation between 280 and 315 nm (UV-B) is strongly affected by the thickness of the ozone layer, whereas a large part of the UV-A radiation (315 – 400 nm) passes through the atmosphere.

³Sometime the wavelength boundaries are listed as 320 and 290 nm (Diffey, 2002).

The intensity of the UV radiation that reaches the surface of the Earth is influenced by many factors:

- The time of day and year. The height of the sun determines the distance the radiation has to travel through the atmosphere; the longer the distance, the more radiation is absorbed, and the lower its intensity. The intensity of UV radiation peaks in the hours around noon during the summer.
- Geographic latitude. In low-latitude locations, the sun is higher in the sky, and the distance the radiation has to travel through the atmosphere is shorter.
- The altitude of a given location. At higher altitudes, the path through the atmosphere is shorter, and more radiation reaches the surface.
- Clouds. Clouds tend to attenuate the radiation. But in some cases, i.e. broken clouds, the intensity of the radiation reaching the surface is actually enhanced.
- Ozone. Ozone is the main absorber of UV-B radiation in the atmosphere. Therefore the intensity of the radiation that reaches the surface is modulated by the thickness of the ozone layer (at approximately 20-25 km altitude). The mean values vary according to location and season, but are also affected by the actual atmospheric conditions.
- Albedo. The amount of radiation that is reflected at the surface (for example by snow or water) also influences its intensity.
- Aerosol. Any aerosol particles present in the atmosphere scatter and absorb radiation, thereby lowering the latter's intensity.

The World Meteorological Organisation (WMO) and the World Health Organisation (WHO) (WHO, 2002) jointly developed the UV index as a tool to inform the general public about the intensity of solar radiation at a given location. (A definition of the UV index as well as additional information on its interpretation are provided in appendix 1). The higher the index, the more UV radiation reaches the surface, and the greater the attendant risk.

KEY MESSAGES

The UV Index is a scale that relates the intensity of solar radiation to the effects on unprotected skin. The actual effects depend on the type of skin but it is generally acknowledged that protection is needed when the index exceeds 3. In many countries the current and/or expected UV index is provided with the weather forecast.

1.2 Artificial sources

1.2.1 Sunbeds and other tanning equipment


Over a century ago, Caucasians looked upon a fair skin as a token of beauty and social achievement (Carter, 2007). As physical activity and outdoor living became increasingly fashionable in the early 20th century, so did the tanned skin that was believed to reflect such a lifestyle. Sunlight was thought to exert a wholesome impact, which was certainly true with regard to the bone disease known as rickets (Hess & Unger, 1921).

When there was not enough exposure to sunlight, light sources imitating the solar spectrum were used (Albert & Ostheimer, 2002). Even in the 1980s, many households owned sun lamps (Health Council of the Netherlands: Committee on UV-appliances, 1986). But from the 1970s, tanning became the primary purpose of exposure to artificial UV sources (Bruggers et al., 1987; Diffey, 1987). This was accompanied by a shift to radiation sources in which the ratio of UV-A/UV-B radiation is relatively greater than that of midday solar radiation. Underpinning this development was the argument that this allowed to build up a tan without the risk of ‘sunburn’ (erythema) (1987). It also boosted a ‘tanning industry’, and with it, commercial tanning salons and tanning devices in wellness centres.

Technical aspects

Modern sunbeds and other tanning lamps available for the general public emit a variety of spectra and intensities. A Norwegian study measured the emissions of 191 sunbeds in 78 tanning facilities. The results are presented in Figure 1. These data demonstrate, as mentioned above, that UV-A radiation is an important part of sunbed emissions and accounts for roughly half of the erythema effective exposure (Nilsen et al., 2012).

Figure 1 Normalized results of the average emission of a sample of Norwegian sunbeds. From left to right in each of the three diagrams: bench, canopy, facial, sun at 35°N (maximum), sun at 60°N (maximum). Normalization to sun exposure at 35°N. The numbers at the top of bars are the 10 minutes



The European standard EN-60335-2-27 limits the erythemal effective irradiance to 0.3 W/m². In actual practice many appliances exceed this recommended irradiance (Tierney et al., 2013).

⁴ Gibraltar and Cyprus are situated at about 35°N, Oslo, Stockholm and Helsinki are situated at about 60°N.

Legal aspects

There is no harmonised European legislation for tanning salons. In Belgium, the competent authority for regulations concerning the operation of tanning salons and the placing on the market of sunbeds in the FPS Economy, SMEs, Self-Employed and Energy:

- Directorate General (DG) Energy is the market surveillance authority responsible for the LVD (Low Voltage Directive);
- DG Quality and Security is the market surveillance authority responsible for the GPSD (General Product Safety Directive);
- DG Economic Inspection is the market surveillance authority responsible for the inspections of tanning salons.

A tanning salon may only be operated if it meets the general safety requirements laid down in the Code of Economic Law (*Code de droit économique*), Book IX on the safety of products and services. The requirements of this Code are supplemented by the specific requirements of the Royal Decree (RD) of 20 June 2002 on conditions regarding the operation of tanning salons, last amended by the Royal Decree of 22 October 2010.

In no area of the sunbeds shall the overall erythema effective irradiance (maximum radiation density) exceed 0.3 W/m².

People with type I skin and those aged under 18 are banned from using sunbeds.

Detailed information on the conditions relating to the commercial operation of tanning salons as well as information booklets written by the FPS Economy are available on the internet⁵.

Sunbeds for private use fall under the "low voltage" regulations, i.e. the Royal Decree of 21 April 2016 regarding the placing on the market of electrical equipment). The sunbeds that are placed on the market shall be safe and their overall erythema effective irradiance (maximum radiation density) shall not exceed 0.3 W/m².

Detailed information on these regulations is available on the internet⁶.

Advertising is regulated by Book VI *Pratiques du marché et protection du consommateur* ("Market Practices and Consumer Protection") of the Code of Economic Law.

Information on these regulations is available on the internet⁷.

The tanning salon sector is continuously monitored by the Economic Inspectorate. During inspections, all the administrative and technical requirements laid down in the Royal Decree on tanning salons are verified.

⁵ <http://economie.fgov.be/fr> > Entreprises & Indépendants > Sécurité des produits et des services > Sécurité des centres de bronzage (http://economie.fgov.be/fr/entreprises/securete_produits_et_services/Zonnecentra/)

⁶ <http://economie.fgov.be/fr> > Entreprises & Indépendants > Sécurité des produits et des services > Sécurité des appareils électriques (http://economie.fgov.be/fr/entreprises/securete_produits_et_services/Securete_des_appareils_electriques/Basse_tension/
http://economie.fgov.be/fr/entreprises/securete_produits_et_services/Securete_des_appareils_electriques/#.VkyJ7tlvdpg)

⁷ <http://economie.fgov.be/fr> > Protection des consommateurs > Pratiques du marché > Publicités et pratiques déloyales (http://economie.fgov.be/fr/consommateurs/Pratiques_commerce/Publicite_pratiques_deloyales/)

In the event of non-compliance, depending on the severity of the infringement and according to the requirements laid down in Book XV of the Code of Economic Law, the measures taken may range from a warning or a settlement submission to temporary seizure of the non-compliant sunbeds or the salon as a whole until all infringements have been removed. When the tanning salon does not meet the given deadline, the case is taken to the public prosecutor's office.

Practice of sunbeds in Belgium

At a national level, the inspection reports from the FPS Economy show that the current legislation is poorly abided by:

- In 2015, 65 tanning salons were inspected. Only 2 salons were managed according to requirements of the Royal Decree, which only amounts to 3.1 % of the inspected premises.
- In 2014, 58 tanning salons and 236 sunbeds were inspected. Only 3 salons were managed in accordance with the requirements of the Royal Decree, i.e. no more than 5.2 %. Of the 236 sunbeds, 12.7 % emitted a radiation intensity in excess of the legal limit of 0.3 W/m². One in four inspected tanning salons accepted individuals under 18 or with skin type I.
- In 2013, 96 tanning salons were inspected. Only seven of them were operated in accordance with the law. Likewise, in 2012 (7.5 %) and 2011 (10.9 %), a minority of salons were run in accordance with the law.
- The most prevalent infringements are:
 - o Lack of information on the risks of artificial tanning;
 - o Lack of control and registration of exposure times;
 - o Missing customer files;
 - o Allowing minors or people with skin type I to expose themselves;
 - o The radiation intensity emitted by the lamps.

Given the fact that tanning salons are regulated by national legislation, it is difficult to compare the degree of compliance to the national laws by member state across the EU.

In 2008-2009, a joint action was set up in several Member States by PROSAFE, a non-profit professional organisation active in market surveillance. A follow-up action was launched in 2010-2011. Belgium participated in these actions.

The results of the first action, called "Sunbeds I" (2008-2009), mainly focused on measuring the erythema effective irradiance (radiation intensity). The participating market surveillance authorities inspected over 300 premises and tested over 500 sunbeds.

Most inspections were carried out at service providers (tanning salons, wellness centres, etc.), and were concerned with safety warnings and advice to consumers, sunbed labelling, the availability of protective eyewear and the measuring of the UV radiation emitted by the sunbeds. The general conclusions drawn from the outcome of the inspections conducted during this first action were the following:

- Tanning salons often failed to provide customer advice, a fact that is difficult for an inspector to verify;
- Sunbed labelling was not up to standard in at least 20 % of cases;
- Of the 84 sunbeds measured, 83.3 % emitted radiation in excess of the 0.3 W/m² limit. It should be noted, however, that these measurements were not performed at random, but targeted sunbeds suspected of exceeding the limit. Across the market as a whole, the percentage of offending sunbeds ranged from less than 3 % in one Member State to over 80 % in several other Member States. Inspections conducted as part of the second action, viz. "Sunbeds II" (2010-2011), were based on the same criteria as the "Sunbeds I" action, but included compliance with the minimum age requirement.

The 12 participating market surveillance authorities inspected over 1,150 premises with sunbeds, most of which were tanning salons and wellness centres. The overall conclusions drawn from this second action were the following:

- Tanning salons failed to provide customer information and advice on a regular basis, a fact that is difficult for an inspector to verify;
- In many tanning salons, the minimum age requirement, viz. no under 18-year-olds, was not abided by;
- Sunbed labelling was not up to standard in at least 20 % of cases. In some participating countries, the labelling failed to meet the requirements in up to 20 %, or even 45 % of cases;
- Over half of the sunbeds in tanning salons emitted a radiation intensity that exceeded the maximum allowed. The radiation emitted by 1.072 sunbeds in total was measured as part of the action. As it turned out, 688 (64 %) of these sunbeds (sometimes significantly) exceeded the limit.

Other considerations

As pointed out above, advertising and the promotion of sunbeds claim that the latter are a good source of vitamin D as well as being useful for pre-tanning before leaving on holiday. Furthermore the sector has also been making a number of health claims (preventive action on diseases such as cancer, diabetes, multiple sclerosis, Alzheimer, etc.) in its advertising (websites, social media).

Prevalence of sunbed use in Belgium

According to data from the Belgian Cancer Foundation (2015a), in 2015, the overall prevalence of sunbed use in Belgium was 14 % (at least 1 tanning session during the past 12 months). 62 % of sunbed users have at least 10 sunbed sessions per year, and 23 % more than 20 sessions per year. This amounts to a total of over 1 million people aged between 15 and 65 having used sunbeds during the past year.

52 % of the Belgian population aged between 15 and 65 have used sunbeds at least once in their lives. This amounts to \pm 3 750 000 individuals aged between 15 and 65. On average, first-time sunbed users are 25 years old, with 38 % starting to use sunbeds before the age of 18. In 2015, 20 % of those aged < 35 had used sunbeds at least once during that year.

6 % of families privately owned a sunbed in 2015. Sunbeds at home are typically used by several members of the family, in 6 % of the cases this involved family members aged under 18 (table 2).

Table 2 Use of privately owned sunbeds by family members (Belgian Cancer foundation, 2015a).

Which family members make use of your sunbed at home? (n = 73)					
	<i>Yourself</i>	<i>Your partner</i>	<i>Children (< 18 years of age)</i>	<i>Children (> 18 years of age)</i>	<i>Other adult family members</i>
<i>Yes</i>	71 (97%)	35 (48%)	4 (6%)	14 (19%)	15 (20%)
<i>No</i>	2 (3%)	19 (26%)	35 (48%)	27 (36%)	26 (36%)
<i>Not applicable</i>	0 (0%)	19 (27%)	34 (46%)	33 (45%)	32 (44%)

The percentage of sunbed users is declining slightly. The percentage of sunbed owners at home has remained stable over the past years.


During a Euromelanoma screening campaign conducted in Europe between 2009 and 2012 (appendix 2) and during which 38 484 people were screened, it was found that the overall average sunbed use in 31 European countries was 8.6 % (all ages, both genders). Belgium turned out to have an above average (15 % on average, gender and age confounded) prevalence of use, i.e. 23 % within the screened population. Focussing on the female population aged under 35, Belgium appears to have the highest rate of sunbed use, viz. around 30 % (less than 20 sessions per year). On average, sunbeds had been used for 8.35 years (median 6). From 2009 to 2014, 18 000 Belgians were screened, and among the total population screened (aged between 15 and 35, both males and females) 25.6 % had used sunbeds at least once, 90 % for less than 20 sessions per year and 10 % for more. These data are not representative for the general population, but the number and the reproducibility of these data made them reliable.

Characteristics and behaviour of sunbed users in Belgium

Positive determinants of sunbed use are: female gender, skin type II, being single (widow or divorced). The positive correlation with other risk factors such as smoking and sunny holiday destinations were measured. There was no overall positive or negative correlation with social class. Still, as it turns out, with 15 % of sunbed users coming from the lowest social class (10.6 % in the Belgian population), the latter is overrepresented (Belgian Cancer Foundation, 2015b).

Reasons for using sunbeds

The main motivation for using sunbeds is a desire for a tanned appearance (getting a tan 62 %, keeping one's tan 44 %) or preparing the skin before a holiday (47 %). 32 % of users indicate that it makes them feel better, 9 % use sunbeds to boost their vitamin D levels.



Knowledge and misconceptions

Though the population is generally well aware that sunbed use is carcinogenic (94 % agree), misconceptions continue to exist. The main misconceptions are:

- Pre-tanning on sunbeds helps to avoid sunburn from the sun: 40 % mistakenly believe this to be the case (44 % believe a tanned skin protects them from the sun).
- A larger percentage of sunbed owners underestimate the risks of sunbeds compared to non-owners. They also have a more positive attitude towards sunbeds:
 - o 18 % of non-owners think using sunbeds is safer than sun-tanning, compared to 47 % of sunbed owners;
 - o 36 % of owners versus 55 % of non-owners think sunbeds should be banned.

KEY MESSAGES

The prevalence of sunbed use in Belgium is high compared to other European countries.

The main reason for using sunbeds is to get a tan (for purely cosmetic purposes).

Sunbed users are predominantly female and young (< 35 years).

1.2.2 Other sources of UV radiation

UV sources other than the sun and tanning lamps are also used for both medical and non-medical purposes. Phototherapy has a one-century-long history (Albert & Ostheimer, 2002, 2003a, b). It started with the idea that using lamps as a means to provide 'sunlight' prevented the occurrence of a variety of common diseases (the so-called Finsen-therapy). Although various treatments have now become obsolete, phototherapy with or without photosensitisers is used to treat a range of skin diseases (Ceburkov & Gollnick, 2000; Diffey, 2002; Choudhary et al., 2009). Both broadband sources and more monochromatic sources are used to induce exposure in the UV and visible spectral region.

In occupational settings a variety of sources are used. An overview is presented in Table 3 (2009). Protection standards in the form of occupational exposure limits have been published by the ICNIRP⁸ (2010) and the ACGIH⁹ (2010).

A more detailed discussion of these sources and exposures would go beyond the scope of this report.

8 International Commission on Non-Ionizing Radiation Protection.

9 American Conference of Governmental and Industrial Hygienists.

Table 3 Common UV sources in the workplace. Derived from Ontario Government (2009).

Source	Potential for Overexposure	Hazard Description
The sun†	Very high	UV from the sun is highest in spring and summer from 11 to 16 hours. UV guidelines can be exceeded in 15 minutes on a clear summer day. Clouds may do little to reduce UV levels.
Electric Welding Arcs	Very high	Welding arcs can exceed the UV guidelines in seconds within a few meters of the arc. Besides workers, bystanders and passers-by are often overexposed to UV from the arcs.
UV Curing Lamps	Medium	Lamps are usually inside cabinets, but substantial UV radiation can escape through openings.
Black Lights	Medium to Low	Low-power UV-A lamps used in non-destructive testing (NDT), insect control, and entertainment.
Germicidal Lamps	High	UV-B- and UV-C-emitting lamps used to sterilise work areas in hospitals and laboratories.
UV Lasers	High	Source of intense UV radiation at a single wavelength, with no visible light.
Lighting	Low	Most lamps used for lighting are made to emit little or no UV radiation.

† The sun has been included for completeness as it is an important source of occupational UV exposure.

2 Effects and health risks of UV

2.1 Intentional effects

2.1.1 Photo-protection effects

The degree of natural pigmentation in the skin rises from phototype II to VI and as the synthesis of eumelanin in the skin increases, so does the protection conferred. The purpose of tanning is to induce facultative non-permanent tanning, which is poorly efficient in providing protection against UV-induced damage. In contrast, the constitutive melanin present in naturally pigmented skin (phototypes IV to VI) has a great ability to protect against skin cancer. There are 6 skin phototypes. People with phototype 1 skin tend to have ginger hair or be albinos. They never tan but always burn when exposed to sunlight. People with phototype 2 skin tend to have blond hair, tan slightly and always burn when exposed. People with phototype 3 skin tend to have blond or light brown hair, get a moderate tan and sometimes burn. People with phototype 4 skin tend to have a Mediterranean appearance with olive skin; they burn very little and build up a deep tan. People with phototype 5 skin exhibit significant pigmentation; they build up a deep tan and burning is rare. They have a Mediterranean, Latin-American or North-African appearance. People with phototype 6 skin are black individuals of Central- or South-African descent; they tan slightly and never burn (Fitzpatrick's Dermatology in General Medicine. McGraw-Hill Professional; 5th edition).

The process that induces pigmentation following UV radiation has been identified. As previously shown, UV radiation triggers the formation of pyrimidine dimers (thus directly resulting in DNA damage). More recently, it has been found that, as a corollary, the dimers generate an intracellular (keratinocyte) increase and activation of the p53 protein. p53 will transactivate the transcription of different genes, pro-opiomelanocortin (POMC), endotheline 1, and many others that are related to the process of UV irradiation (inflammation, pigmentation, etc.) (Chen et al., 2014).


The fact that pyrimidine dimers are the direct inducers of pigmentation is proof that skin pigmentation is only related to DNA damage.

When POMC is synthesised, posttranslational cleavage of POMC produces β -endorphin and α -MSH (melanocyte-stimulating hormone). Beta-endorphin synthesis may be responsible for tanning addictions.

Once the MSH-receptor (MC1R) is activated by its ligand (MSH), the synthesis of eumelanin (black and brown melanin) is favoured instead of pheomelanin (red). Eumelanin provides protection against UV damage because it is mainly a free radical scavenger and acts as a chromophore. In contrast, UV radiation causes pheomelanin to be a potential source of free radicals that are also responsible for oxidative DNA damage.

Both types of melanin are synthesised inside specific organelles, called melanosomes. Melanosomes are transferred from the melanocytes to the keratinocytes and confer a pigmentation to the epidermis. Skin pigmentation is related to the amount and the type of melanin synthesised (pheomelanin (red) and eumelanin (brown to black)).

Uv mediated skin tanning



• Tran T-N et al, *Pigm Cell Melanoma Res*, 2008

Given the fact that UV-induced pigmentation provides low photoprotection (equivalent to a sunscreen with an SPF-level of 2), UV protection is mandatory, especially for children and fair-skinned individuals (even lower photoprotection) (Miyamura et al., 2011). Additionally, these authors were able to confirm that UV-A tanning does not confer any photoprotection but does induce DNA damage, just as is the case for UV-B rays. Many people mistakenly believe that a few artificial UV sessions prior to sunny holidays may protect them against sunburns whilst in fact, the low level of photoprotection induced by UV tanning does not prevent the occurrence of sunburns and UV-induced DNA-damage (Brenner & Hearing, 2008).

KEY MESSAGES

UV-induced skin pigmentation is only obtained as a result of the formation of pyrimidine dimers (i.e.: DNA damage).

Constitutional skin pigmentation (phototypes IV to VI) provides protection against skin cancer but UV-induced pigmentation will only provide protection to the skin in the range of an SPF-level of 2-4 (equivalent to sunscreens with an SPF of 2).

UV-induced pigmentation confers low photoprotection. Artificial UV exposure prior to sun exposure should be avoided. There is therefore no good reason to make use of artificial UV-exposure as a means to prevent sunburns.

2.1.2 Sociological and psychological effects

Sociological aspects

Mankind has always had a special and privileged relationship with the Sun. Both worshipped and feared, many supernatural powers have been attributed to this celestial body, which has always had a prominent place in our civilisations.

It should be noted, however, that depending on the time and/or place, it has triggered quite different types of behaviour.

A few examples give a good account of sociological data grounded in habits and customs that may fluctuate according to fashion trends.

In this part of the world, a fair skin used to be associated with a form of nobility, whereas a darker skin tone hinted at peasantry. This has been a thing of the past for several decades now.

The history of sun tanning reveals that attitudes started to change around 1920-1930. When paid leave became a reality in 1936, workers started to travel and seek destinations with plenty of sunshine. After World War II, as the bikini became fashionable, exposure to the sun became all the more popular. Thus, it took no more than a fashion trend to allow these bodies to tan in the sun.

A tanned appearance has become indicator of "good health" (cf. "a healthy tan"), whereas a white or even pallid skin has clearly grown to be interpreted as a sign of illness. Advertising has been eager to shape consumer behaviour accordingly.

After sunscreens, self-tanners appeared on the market. Next came sunbeds as a means to extend this pursuit of sunshine, regardless of the weather and the location.

Psychological aspects

Besides this sociological aspect, the psychological factors involved in the decision to use a sunbed also need to be mentioned. What is the basis for the desire to go to a tanning salon or the irresistible urge to return frequently? Are some people more vulnerable than others? When does a type of addiction, i.e. dependence, set in?

Psychologically, the starting point is self-image. This self-image is linked to one's "outer persona". It is a "calling card" that is shaped during the first years of life and allows one to position oneself in the eyes of others as in our own.

This self-image tends towards a certain search for an ideal, whilst being moulded by the cultural and social context. This self-image goes beyond our earthly bodies in the sense that it also involves psychological components that may or may not be consistent with the representation of the body. This is why everyone has a physical and psychological image of themselves that may fluctuate in response to the events of life. Moreover, this self-image faces the dictates of fashion. Over the last few decades, fashion has imposed its laws, bringing in its wake conformism and anti-conformism. Tanning is an adornment, just as clothing is. This reflects the growing importance of the representation of the body in our societies. Despite some reactions to tanning, as was the case in the 1990s when Calvin Klein and Kate Moss advocated pale complexions, a fashionably tanned skin remains synonymous with good health and, for some, a high social standing.

Behind the fashion trends there looms, at the psychological level, a whole narcissistic dynamic. The culture we live in is admittedly increasingly narcissistic and great value is attached to surpassing oneself (e.g. sporting competitions), whereas illness and death, if not simply ignored, give rise to a great deal of fear.

It follows that, within such a cultural context, displaying some degree of narcissism is normal. Still, calling someone narcissistic remains pejorative. Most notably, such a person's behaviour is characterised by a lack of psychological depth as well as superficiality in their interaction with other people. This accounts for the fact that there is still an ambiguous and ambivalent connotation to narcissism.

There does not seem to be any real at-risk type of personality, but some types are clearly more vulnerable to narcissistic injuries.

Thus, when the personality traits grow rigid, inadequate, persistent and significantly impair functioning or cause subjective suffering, this will be consistent with a personality disorder, i.e. a predisposition.

Thus, according to the DSM-IV, a "narcissistic personality disorder" is characterised by a pervasive pattern of grandiosity, need for admiration and lack of empathy.

Recently, Fell et al. (2014) emphasised how dangerous exposure to the sun can be, not only as regards the risk of cancer, but also the risk of addiction. Chronic exposure to UV radiation releases endorphins in the body that elicit the same chemical response as opiates, e.g. heroin or morphine. These are the same receptors that could cause symptoms that are typical of intoxication, which is consistent with the findings of laboratory research with mice (Petit et al., 2014). This compulsive desire to build up a tan, regardless of the proven negative consequences, is suggestive of a behavioural complex similar to addictive disorders (Warthan et al., 2005; Kourosh et al., 2010; Robinson et al., 2010; Harrington et al., 2011; Hillhouse et al., 2012).

It is believed that such an addiction most frequently affects adolescents (Petit et al., 2011, 2013, 2014). Depending on the parameters used, various studies take the view that 5 % to 15 % of sunbed users can be considered to have a tanning addiction (Heckman et al., 2014). This addiction mainly affects young girls, but is also found in the least-favoured social classes. The label "tan-addict" has been proposed to describe an irrepressible urge to expose oneself to UV radiation, bringing about a sense of lack when this activity cannot be engaged in and a feeling of euphoria and an enhanced mood when it can, as well as a relaxing and anxiolytic effect. Moreover, knowing what the adverse health effects of this type of behaviour are does not prevent it from continuing.

This diagnosis of "tan-addict" has led to various debatable translations in French, viz. "*tanorexie*, *tanoholisme* or *bronzomanie*" (WIKIPEDIA), i.e. many neologisms to describe the pathological addiction to tanning. Indeed, it is necessary to understand the nature of the addiction and pathological need to have a tanned skin underlying all these pseudo-diagnoses. The quest for an aesthetic appearance and social desirability has been described as the main factor that accounts for the fact that health concerns fade in the background.

This addiction may occur in combination with other psychiatric disorders. Thus, an American study (Mosher & Danoff-Burg, 2010) showed that out of a total of 229 students, 70 (30.6 %) of those who had a pathological addiction to tanning based on the diagnostic criteria of DSM-IV-TR and CAGE, also reported anxiety symptoms and an excessive use of alcohol, marijuana and other substances. In another study (Ashrafioun & Bonar, 2014) involving a sample of 533 students, 31 % had a tanning addiction. This tanning addiction was predominantly found in females, and there was a significant association with an obsessive-compulsive disorder as well as a body dysmorphic disorder.

KEY MESSAGES

The sociological data account for changes in habits and customs as regards the relationship mankind has had with the sun, which varies from one period in time to another, and from one continent to another, and is significantly affected by the dictates of fashion.

In this part of the world, tanning started to become fashionable in the 1920s-1930s, a trend that gained momentum as paid leave and lengthier holidays became a reality, with a tanned appearance becoming synonymous with "good health". Fed by the media, this quest for sunshine naturally resulted in the emergence of sunbeds.

On a more psychological level, this desire for a tanned appearance is grounded in the image of oneself one wishes to convey to others, and which is linked to one's "outer persona" and the search for a certain ideal. Whilst this drive to attract and seduce typically finds its roots in the narcissistic aspects of our personalities. It can, however, result in a form of addiction in some more vulnerable individuals. Based on the release of endorphins during chronic UV exposure, this pathological addiction to tanning, known as tanorexia, is mainly found in adolescents, mainly female, often in association with other psychiatric disorders such as anxiety disorders, obsessive-compulsive disorders, body dysmorphic disorders and addictive disorders.

2.1.3 Vitamin D

In 2016, the SHC of Belgium issued the sixth edition of the “Nutritional recommendations for Belgium” (SHC 9285, 2016). For each nutrient including vitamin D, the new recommendations provide the following information:

- The nutritional properties and the impact on health,
- The recommended intake for different age/gender groups,
- The risks related to an excessive intake and the estimated upper tolerable intake,
- The alimentary sources and the average intake by the Belgian population,
- Practical recommendations (incl. dietary supplements),
- References.

Synthesis

Vitamin D is an essential hormone that has a considerable influence on the calcium balance and bone development.

Humans acquire vitamin D through endogenous or exogenous synthesis. Endogenous synthesis of vitamin D₃ occurs in the skin upon exposure to UV-B radiation. Exogenous vitamin D₂ or D₃ is supplied through dietary intake, mainly through oily fish (e.g., salmon), dairy products, mushrooms and eggs. Many food products are nowadays also enriched with vitamin D₃.

Vitamin D on its own has no physiological action. To be physiologically active, vitamin D must undergo two chemical transformations. The first transformation occurs in the liver, which yields 25-hydroxyvitamin D. The second takes place in the kidneys, yielding 1 α , 25-dihydroxyvitamin D. It is under this form that vitamin D is physiologically active. Because serum 25-hydroxyvitamin D levels are affected by the intake of vitamin D as well as the exposure to UV-B rays, measuring serum 25-hydroxyvitamin D levels is commonly used as a marker of the “individual vitamin D status”.

In the early 2000s, it was discovered that many tissues other than the kidney could locally produce 1 α , 25-dihydroxyvitamin D. This discovery gave rise to the hypothesis that active vitamin D produced at tissue level could play a role in the healthy functioning of organs.

UV-lamps and endogenous vitamin D synthesis

Vitamin D is only synthesised to a limited extent in the skin, and only during the initial exposures to UV radiation. Moreover, this process is largely dependent upon the intensity of the UV-B radiation emitted by the lamps (Davies et al., 1980; Sallander et al., 2013).

Two controlled interventional studies have been conducted on the influence of typical sunbed exposures on serum 25-hydroxyvitamin D levels.

A Danish group performed a randomised trial that involved Caucasian females aged 50 years and over, assigned to:

- (a) A control group (21 women),
- (b) A group (n=20) undergoing 4 UV-A-tanning sessions for 6 to 12 minutes on machines emitting 0.4 % UV-B rays,
- (c) And a group (n=15) undergoing 4 UV-A-tanning sessions on machines emitting 1.4 % UV-B rays.

As it turned out, 37 to 64 % of sunbed sessions induced side effects such as erythema and polymorphic light eruption. The average baseline serum 25-hydroxyvitamin D level was 19 ng/mL. 25-hydroxyvitamin D levels did not change in the control group. After 4 sunbed sessions, they significantly increased by an average of 12.5 nmol/L and by 27.5 nmol/L in the 0.4 % and 1.4 % UV-B groups, respectively. After 4 more sessions, these increases became non-significant, viz. only 3 and 0.5 nmol/L, respectively. Thus, a plateau was rapidly reached after only a few sessions as regards the circulating 25-hydroxyvitamin D levels (Thieden et al., 2008).

Another Danish group randomised 20 healthy volunteers to one 10-16 minute whole-body exposure using either a commercial sunbed with UV-B emissions (UV-B/UV-A ratio 1.8 – 2.0 %) or an identical placebo sunbed that did not emit UV-B rays (Langdahl et al., 2012). After seven days, the mean increase in 25-hydroxyvitamin D levels in the UV-B group was 4.5 nmol/l (SD 7 nmol/l) compared to a decline of -1.2 nmol/l (SD 7 nmol/l) in the placebo group ($p = 0.1$).

Two trials have tested the influence of extensive (10 minutes or more) sunbed exposure sessions over a short period of time on the vitamin D status.

In the Netherlands, a trial was conducted with 105 young adults (aged between 18–30 years; 91 % female) during the winter months. 35 of these adults were randomly assigned to a group receiving 6-to-12 minute exposures to sunbeds in commercial indoor tanning salons 3 times a week for a period of 8 weeks. Hence these subjects were exposed to a total of 24 tanning sessions (de Gruijl et al., 2012). Another thirty-seven subjects received 25 µg per day of vitamin D supplementation, and the remaining 33 subjects were assigned to a control group who received neither vitamin D supplementation nor sunbed sessions. After 8 weeks, the mean serum 25-hydroxyvitamin D levels dropped from 62 to 55 nmol/L (-7, SD:11) in the control group. They rose from 62 to 109 (+47, SD: 23) and from 58 to 93 nmol/L (+35, SD: 19) in the sunbed and in the supplementation groups, respectively. The statistical difference between the sunbed and the supplementation groups was not reported.

A Norwegian randomised cross-over trial involved 31 healthy volunteers (8 men and 23 women) aged 21 to 61 who, during a 2-week period, either underwent a series of ten commercial sunbed sessions for 7 to 15 minutes (equivalent to a mean of 12.6 SED) or received a daily dose of 50 µg vitamin D (Lagunova et al., 2013).

Nine subjects withdrew from the trial, leaving 11 subjects in each group. A similar increase in 25-hydroxyvitamin D levels was observed in both groups, i.e., 19•8 (SD: 5•4) nmol/L after the sunbed sessions and 25•3 (SD: 5•4) nmol/L after supplementation (Lagunova et al., 2013).

Hence, in fair-skinned people, short sessions (a few minutes) of UV-B irradiation of the skin can trigger a significant increase in 25-hydroxyvitamin D levels and longer exposure times do not enhance endogenous vitamin D production. Sunbed sessions slightly increase these rum 25-hydroxyvitamin D levels, but only during the first sessions. The trials that involved more sunbed sessions suggest that in healthy adults, a large number of sunbed sessions with short intervals (2-3 days) in between does not boost the vitamin D status to levels beyond what daily supplementation with oral vitamin D would achieve.

Health effects of vitamin D

It has been known for about a century that the physiologically active form of vitamin D is necessary for the digestive absorption of calcium and phosphorus in the gut and its incorporation in bone tissues. Childhood and pregnancy are two periods of life during which vitamin D requirements are higher.

Ecological, cross-sectional and observational studies have documented the association between a low vitamin D status and a greater risk of being diagnosed with a myriad of diseases, including minor ailments and rare conditions (Harvey et al., 2012). Many have concluded from these non-interventional studies that a low vitamin D status could be a cause of illness/disease, and that preserving a high vitamin D status could provide an easy way to prevent diseases and increase life expectancy. In the same spirit, it was believed that maintaining high serum levels of circulating 25-hydroxyvitamin D would allow the optimal local production of physiologically active vitamin D at tissue level.

A more recent study looked into the different beneficial health outcomes initiated by UV radiation through Vitamin D-dependent and independent pathways such as the beneficial effects on infectious diseases, inflammatory skin diseases (e.g. psoriasis), asthma, cardiovascular diseases, obesity and diabetes and all-cause mortality (Hart et al., 2017).

Vitamin D is in fact a naturally occurring hormonal drug. As for any drug, the most robust way to evaluate the health effects of increasing vitamin D levels is to examine how increases in circulating 25-hydroxyvitamin D impact on the incidence and mortality associated with conditions deemed to be vitamin D-preventable.

To date, the over 400 randomised trials that have been conducted on vitamin D supplementation and their meta-analyses have been unable to confirm the health benefits supposedly associated with increased 25(OH)D levels, even when high doses of supplementation (i.e., ≥ 50 μg per day) were used in subjects with a low vitamin D status before randomisation, and even on skeletal or postural endpoints (i.e., osteoporosis, fractures and falls) (Autier et al., 2014; Bolland et al., 2014a; Bolland et al., 2014b; Bolland et al., 2014c; Meyer et al., 2015; Theodoratou et al., 2014). The risk of all-cause death among elderly subjects is the only outcome for which a significant risk reduction of 11 % has been observed with ordinary doses (5 to 20 μg or 100 to 800 IU per day) of vitamin D₃ supplementation tested in 14 trials (Chowdhury et al., 2014). This effect is not observed with vitamin D₂ supplementation.

However, a more careful reading of the literature shows that if the analysis is restricted to the 10 trials that actually tested vitamin D3 (cholecalciferol) on its own (and not another vitamin D compound [e.g., calcitriol or 1 α ,25-dihydroxyvitamin D3]), the risk of death was reduced by 8 %, which was no longer statistically significant (Bolland et al., 2014). In any case, these results are not relevant to the issue of artificial UV tanning by adolescents and young adults.

Extensive randomised trials are being conducted on vitamin D supplementation (Manson et al., 2015), but the first results that have been released so far do not differ from those of earlier, smaller trials (Amrein et al., 2014). So, the fact that administering vitamin D does not have any impact on virtually any outcomes strongly indicates that low 25(OH)D should be looked upon as a marker rather than a cause of impaired health. The main working hypothesis for explaining the low vitamin D status associated with so many conditions is based on acute and low-grade chronic inflammatory processes that are part of many physiological phenomena involved in deteriorating health and diseases (Autier et al., 2014).

Vitamin D recommended intake

The recommended intake of vitamin D is actually still debated amongst scientists. However, there is at present no firm evidence in support of advising healthy subjects to increase their vitamin D status (Meyer et al., 2015).

The SHC of Belgium recommends a daily intake of 10 μ g Vitamin D3 for children (0 - 10 years), 10-15 μ g for adolescents and adults (11-70 years), 20 μ g for pregnant and breastfeeding woman and 20 μ g for the elderly (> 70 years) (SHC 9285, 2016).

Because of the numerous health hazards associated with exposure to artificial UV radiation (e.g., increased risk of skin and eye cancer, skin rashes, sunburn, skin ageing), individuals who wish to increase their vitamin D status may turn to oral vitamin D supplementation. There is some debate as to the tolerable upper intake level of vitamin D. The SHC of Belgium has set the upper limit at 25 μ g vitamin D3/day for children 1-10 years old and at 50 μ g/day for subjects aged 11 years and older. The cost of vitamin D supplements is generally lower than that of a series of UV tanning sessions and does not trigger the undesirable effects of exposure to artificial UV radiation.

KEY MESSAGES

There is currently no evidence that increasing the vitamin D status of healthy subjects with a normal diet is associated with any health benefits in terms of preventing diseases or improving their outcome.

Exposure to artificial UV radiation as a means to increase the vitamin D status is no more efficient than taking oral supplements of vitamin D, but it is associated with an increased risk of several health conditions, including skin and eye cancer.

People who wish to increase their vitamin D status can do so by consuming oral vitamin D supplements. Oral supplementation with ordinary doses as recommended by the SHC (10 to 20 μ g/day) is safe.

2.1.4 Other uses

The use of UV radiation for therapeutic purposes goes beyond the scope of this advisory report.

2.2 Non-intentional effects

2.2.1 Carcinogenicity

Genotoxicity

Different wavelengths of UV light induce different types of DNA damage (SCHEER, 2016). UV-C and UV-B, but much less so UV-A, are capable of exciting the DNA molecule directly and subsequently generating DNA photoproducts. Indeed, DNA is regarded as a major chromophore for most of the biological effects of UV-B and UV-C, including erythema, tanning, immunosuppression, mutagenesis and carcinogenesis. But other chromophores have also been shown to be involved. Thus, double stranded snRNA (through LTR3), and tryptophan (through AhR), lipids and receptors in the cell membrane contribute to the appearance of erythema, pigmentation and/or immunosuppression. Also, urocanic acid plays a part in immunosuppression, whilst a UV-A-sensitive opsin in melanocytes is involved in immediate pigment darkening. The main UV-induced DNA photoproducts are dimers, which are formed by the covalent binding of two adjacent pyrimidines in the same polynucleotide chain. The two major types of pyrimidine dimers are cyclobutane dimers and 6.4-photoproducts.

While UV-B-induced pyrimidine dimer formation results from the direct absorption of photons by DNA bases, UV radiation (and violet/blue light) can also damage DNA indirectly. As a consequence of photon absorption by non-DNA chromophores, energy can be transferred either to DNA (type I photosensitized reaction) or to molecular oxygen, with reactive oxygen species in turn being able to damage DNA (type II photosensitized reaction).

It is noteworthy that much of the biological impact of UV-A, including its cell toxicity, is largely dependent upon the presence of molecular oxygen, which is indicative of the prominent role played by reactive oxygen species. Although these reactive oxygen species are also formed by UV-B, UV-A has been shown to be largely responsible for the oxidative DNA damage that occurs after the exposure of cells to natural sunlight. UV-A-induced pyrimidine dimer formation may involve a type I photosensitized reaction, unlike direct UV-B induced pyrimidine dimer formation.

UV-induced reactive oxygen species include singlet oxygen and probably other non-radical and radical reactive oxygen species, such as hydrogen peroxide and the superoxide radical. This oxidative stress affects not only DNA, but also membranes and proteins. The relative contribution of each (oxidative membrane damage, oxidative protein damage, oxidative DNA damage) to the different biological effects of UV irradiation has not been well established.

Mutagenesis and carcinogenesis

The wavelength dependencies (the 'action spectra') of pyrimidine dimer formation, cell killing, mutation, and cell transformation are highly similar and resemble the DNA absorption spectrum (at least over the UV-C and UV-B ranges, but also up to 330 nm into the UV-A range). Both cyclobutane pyrimidine dimers and (6-4) photoproducts have been shown to be mutagenic, causing typical 'UV-signature' mutations, i.e. C→T transitions at neighbouring pyrimidines in the same DNA strand (sometimes even tandem CC→TT mutations). Skin carcinomas show these UV-signature mutations abundant in the p53 tumour suppressor gene, which constitutes classical evidence that UV radiation plays a part in triggering these skin cancers (SCHEER, 2016).

Next Generation Sequencing techniques have revealed that - despite efficient DNA repair mechanisms - sun-exposed middle-aged skin is littered with mutations, with 15 % of the epidermal cells carrying a p53-mutation, most of which bear the UV signature. A recent study conducted on 74 cancer genes found that there were over 100 mutated cell clones per cm² of eyelid skin from people over 55 who had undergone eyelid surgery, and again most of these mutations bore the UV signature.

Given these findings, it is not surprising that skin cancers rank among the absolute top in terms of the number of mutations they carry (10 000 to over 100 000 per genome, with hundreds in coding regions). And again UV-signature mutations predominate. Due to this wealth of mutations it is difficult to separate the 'driver' mutations from the 'passenger' mutations, i.e. determine which mutations really drive tumour development and which mutations are just hitchhiking.

But before NGS was available, driver mutations had already been identified by studying familial/hereditary skin cancers (e.g., p16Ink4A loss in FAMMM melanomas, and defective PTCH activating the Hedgehog pathway in BCC in Gorlin syndrome) and recurrent mutations (e.g. in mutated p53 in skin carcinomas).

Although UV-signature mutations appear to abound in skin cancers, this is not always clearly the case in prominent driver mutations (e.g. PTCH or SMO in BCC, or N-RAS or B-RAF in melanomas), with the exception of P53 in skin carcinomas and Notch1- 4 in squamous cell carcinomas of the skin.

Experiments with transgenic mice carrying one of the prominent driver mutations show however, that UV irradiation can greatly enhance the development of the corresponding skin cancer (Noonan et al, 2001; Noonan et al. 2010; Wolnicka-Glubisz A et al., 2006; Berking et al. 2002)

KEY MESSAGE

Experiments with transgenic mice carrying one of the prominent driver mutations show that UV irradiation can greatly enhance the development of the corresponding skin cancer.

Epidemiological studies (sunbed-related)

After World War II, the incidence of cutaneous melanoma increased sharply in most fair-skinned populations and in the 2000s, the rise in incidence is still a fact in most fair-skinned communities (Erdmann et al., 2013).

A vast body of epidemiological data indicate that the increased incidence of melanoma is mostly attributable to the fact that sun exposure became fashionable and spread during the second half of the twentieth century (IARC, 1992). That the ultraviolet (UV) radiation component of the sun spectrum reaching the earth's surface is the carcinogenic agent was formally recognised more recently (IARC working group, 2006; "Radiation", 2012). Intermittent exposure to UV-sources is believed to be the real culprit in the occurrence of melanoma, intermittency being defined as the brisk exposure of skin areas that are never exposed to sunlight most of the year. Experiments in animals and epidemiological studies (e.g., in migrants) have provided ample evidence that susceptibility to UV is greater at a younger age (mainly in childhood) than later in life (Autier et al., 1998; Noonan et al., 2001; Noonan et al., 2012; Whiteman et al., 2001).

In twenty years' time, sunbed use has become widespread in fair-skinned populations, mainly among young (i.e., under the age of 25) adults, especially women. Sunbed use is a typical intermittent sun exposure pattern.

In large powerful tanning units, the UV intensity may be 10 to 15 times higher than that of the midday sun (Gerber et al., 2002), and the UV-A doses per unit of time received by the skin during a typical sunbed session are well above those experienced during daily activities or sunbathing.

The annual UV-A doses received by frequent indoor tanners may be 1.2 to 4.7 times those received from the sun, and these doses add to those received from the sun (Miller et al., 1998). However, the UV emission spectra of sunbeds have special characteristics and are different from the solar spectrum: sunbed emission spectra are similar to the sun spectrum in the UV-B (280-320 nm) range but reach values 10 to 15 times higher in the UV-A (320-400 nm) range (Gerber et al., 2002). The average UV output of sunbeds corresponds to a UV index of 13, which is what vacationers encounter on Mediterranean beaches in the summer. In sunbeds, the trunk is the most exposed area of the body, whilst the head and feet are the least exposed, though there are sunbeds that are fitted with high-pressure lamps specially designed for the face. Natural sources of UV-A are probably never this powerful and repeated exposures to high doses of UV-A constitute a new phenomenon in human beings.

For obvious ethical and practical reasons, there are no direct experimental human data available (e.g., a randomised trial) showing that exposure to sunbeds causes melanoma or other skin cancers. Despite this, in 2009, an international group of experts convened by the IARC classified exposure to "UV-A-lamps" or "UV-emitting tanning devices as carcinogenic to humans" (El Ghissassi et al., 2009). This classification is based on the accumulation of compelling evidence that sunbed use can be a cause of melanoma and not just a proxy for sun exposure for the following reasons:

- Basic animal and human studies have demonstrated that the UV-A and UV-B regions of the radiation spectrum emitted by sunbeds are carcinogenic agents to humans (El Ghissassi et al., 2009; "Radiation," 2012);
- Meta-analyses of case-control and prospective studies have revealed the following:
 - o Based on dose-response calculations, 35 studies provide evidence of a 1.8 % (95 % confidence interval 0 % to 3.8 %) increase in the risk of melanoma for each additional session of sunbed use per year (Boniol et al., 2012).
 - o Three studies show that sunbed use may be involved in the occurrence of eye melanoma (El Ghissassi et al., 2009; "Radiation", 2012), a type of cancer that occurs more rarely than cutaneous melanoma, but is much more fatal.
 - o In 13 studies, first use of sunbeds before the age of 30 is associated with the occurrence of cutaneous melanoma, eye melanoma, basal cell carcinoma and squamous cell carcinoma (Boniol et al., 2012; "Radiation", 2012; Wehner et al., 2012). This relationship has been consistently found in all observational studies.
- Most epidemiological studies on sunbed use and skin cancer have adjusted for sun exposure and natural susceptibility to UV.
- The investigation of a melanoma epidemic in Iceland (a country located between 64 ° and 66 ° N and where sunny days are uncommon) revealed that after 1990, the incidence of melanoma increased sharply, mainly in young women, with preferential occurrence on the trunk. The incidence tended to decline after 2000, when public health authorities imposed more stringent controls over sunbed facilities and their use. Although this was an ecological study, the exposure of Icelandic youngsters after 1985 seemed to be the most likely cause for that epidemic (Alberg, 2011; Autier, 2010).
- Before 1990, the increased incidence of melanoma in women mainly concerned the lower limbs, whereas since 1990, this increase has been more marked on the trunk than on other body parts, which is believed to be the consequence of artificial UV tanning.

The relationship between sunbed use and melanoma that has been observed both in the laboratory and in epidemiological studies is consistent with the data linking sun exposure to melanoma, notably the greater susceptibility to the carcinogenic action of UV radiation at younger ages (Autier et al., 2008; "UV radiation & skin cancer. The science behind age restrictions for tanning beds", 2012). Moreover, epidemiological data on sunscreens, sunbeds and other factors support the findings from basic scientific research that has revealed the carcinogenic properties of UV-A radiation (Agar et al., 2004; Autier et al., 2011; Mouret et al., 2006; Noonan et al., 2012).

The high levels of sunbed exposure in fair-skinned communities will contribute to the continuing upward trend in the incidence of cutaneous melanoma and of other skin cancers (Boniol et al., 2012; Tierney et al., 2015).

Health risks

In 2012, a meta-analysis and review based on 13 informative studies led to the conclusion that first-ever use of sunbeds before the age of 35 was associated with a significant increase in the relative risk of melanoma of 1.87.

By using prevalence data from surveys and data from Globocan (2008) (for 18 EU countries), an estimated 3 438 cases of melanoma could be attributed to sunbed use, most of which affected women (n=2.341) (Boniol et al., 2012).

A similar meta-analysis has been performed on NMSC and included 12 studies with 9 328 cases of NMSC. As regards people who reported ever using indoor tanning compared with those who had never done so, the summary relative risk for squamous cell carcinoma was 1.67 and that for basal cell carcinoma was 1.29, though the latter was not significant.

In the US, the population attributable risk fraction was estimated to be 8.2 % for squamous cell carcinoma and 3.7 % for basal cell carcinoma: this corresponds to more than 170 000 cases of NMSC each year that can be ascribed to indoor tanning. This risk is higher if sunbed use occurred early in life (< 25 years) (Wehner et al., 2012).

KEY MESSAGES

UV-A and UV-B are both carcinogenic.

Recent reviews of biological data, animal experiments and meta-analyses of the risks involved in UVR exposure (including sunbed use) confirm that there is an association between exposure to natural and sunbed UVR on the one hand and the risk of skin cancer, especially melanoma, as well as squamous cell carcinoma.

Economic burden of skin cancer in Belgium

A bottom-up cost-of-illness study was performed for skin cancer in 2015 by the University Ghent (Pil et al., 2016). The total annual cost that is currently induced by skin cancer in Belgium is estimated to be € 103 million (for a population of 8.8 million adults), of which almost € 64 million are borne by the healthcare system (government), which amounts to about 0.15 % of the total healthcare budget in Belgium. Projections to 2034 showed an estimated annual discounted cost of € 153 million, and a total cumulative cost of € 3.2 billion.

A Markov model simulation over 50 years estimated that an average € 227.7 million (0.57 %) of the healthcare budget could be redirected to other diseases by implementing a skin cancer prevention campaign and a total of € 238 million could be redirected by banning sunbeds in Belgium (cumulative costs over 50 years). When calculating the cost-effectiveness of both interventions, they are cost-saving on the long term and thus to be preferred. Over a period of 50 years, 5 944 deaths could be avoided by means of an annual UV prevention campaign (2 368 in males and 3 576 in females) and 5 692 by means of a ban on public sunbed use (2 198 in males and 3 494 in females). Although the fact that over this 50 years period, other factors might also influence the incidence of melanoma in Belgium, this study of Pil et al indicates that a nation-wide population-based strategy promoting UV protective behaviour combined with a national ban on the use of sunbeds can have a positive effect on both public health and public finances (Pil et al., 2016).

KEY MESSAGE

Estimates of the effects of a skin cancer prevention campaign combined with a total ban on sunbed use predict that this would result in the prevention of nearly 6 000 melanoma deaths over 50 years and that € 227.7 million could be saved by a sensitisation campaign and € 238 million by a total ban on sunbeds in Belgium.

2.2.2 Ageing (heliodermy)

Intrinsic skin ageing is a natural process that is determined by genetic and chronological factors that are affected by external factors such as the sun or sunbeds.

Also, the genetic material of the cells is altered, especially the structure of DNA, under the influence of UV-A radiation in particular. In addition, UV radiation triggers the formation of aggressive free radicals, which also modify the structure of DNA. There are physiological DNA repair mechanisms, but these processes are overwhelmed in the event of significant repeated and cumulative doses of UV radiation, causing the skin to change slowly over the years, a process known as “**actinic elastosis**” (Calderone DC et al., 1995).

Excessive sunlight on the neck and cleavage causes the skin to become less supple and atrophic, with increased and decreased pigmentations producing an irreversible appearance called **poikiloderma**.

With brown spots, **solar lentigines**, or depigmented spots called **guttate hypomelanosis** (Nakamura et al., 2015; Ortonne et al., 1990).

Actinic keratoses are precancerous lesions, some of which progress into squamous cell carcinoma, viz. between 0.1 and 0.53 % per year or 5 to 20 % within 10 to 20 years (Trakatelle M et al., 2016; Green AC et al., 2015)

KEY MESSAGES

Heliodermy is the extrinsic ageing of the skin induced by intense and/or frequent and cumulative exposure to the sun and is linked to the harmful effects of ultraviolet radiation, including sunbeds.

There are physiological cellular DNA repair mechanisms, but these repair processes are overwhelmed in the event of significant repeated and cumulative doses of UV radiation, causing the skin to change slowly over the years to progress towards a precancerous state called a “**dysplastic state**”.

Actinic keratoses are UV-induced precancerous lesions that can progress into a squamous cell carcinoma within 10 to 20 years in about 5 to 20 % of cases.

2.2.3 Other issues

Ocular problems linked to UV radiation

Exposure to sunbeds could also increase the risk of eye (intra-ocular) melanoma, a rare tumour that is deadlier than skin melanoma. Four case-control studies conducted so far on artificial UV and intraocular melanoma have all obtained results that are highly consistent with a raised risk: in the US, Tucker et al. 1985 found a 1.4-fold increased risk (95 % CI: 0.9; 2.2, Holly et al. 1990 a 3.6-fold increased risk (95 % CI: 1.6-8.7), and Seddon et al. 1990 a 3.4-fold increased risk (95 % CI: 1.1-10.3) whereas in Australia, Vajdic et al. 2004 found a 1.7-fold increased risk (95 % CI: 1.0-2.8) (Holly et al., 1990; Seddon et al., 1990; Tucker et al., 1986; Vajdic et al., 2002).

Photoallergic reactions

Photoallergic reactions are abnormal reactions induced by an allergy to one or more types of solar radiation – UV-B and/or UV-A and/or visible light - and take on different shapes (Onoue S. et al., 2017).

These reactions are rare and occur in predisposed individuals with a history of allergy. They appear after a certain time of exposure and affect exposed areas of the skin, but may spread to unexposed areas. A short enumeration of the different types of photoallergic reactions is mentioned below. A more detailed description of these diseases goes beyond the scope of this report.

- a) Seasonal episodes of light eruption (Gruber-Wackernagel et al., 2014; Lava et al., 2013):
 - Benign summer light eruption.
 - Juvenile spring eruption.
 - Benign winter light eruption.
 - Polymorphic light eruption.
- b) Chronic actinic dermatitis (Rodríguez-Carreón et al., 2015).
- c) Solar urticaria (Goetze et al., 2015).

Photoinduced disorders

Some intrinsic skin diseases may worsen under the influence of UV radiation. Examples are Lupus erythematosus (Kuhn et al., 2005) and Porphyrias (Palma-Carlos et al., 2005).

Photosensitisation reactions

Photosensitisation reactions are common and can affect anyone from the first exposure to UV radiation, both natural and artificial. They are triggered by the interaction between UV radiation and a toxic substance found in the skin to a greater or lesser extent.

Endogenous photosensitisation occurs when this substance has been ingested. Exogenous photosensitisation occurs when it has been applied to the skin. In this case, the disorder may be either due to a phototoxic or a photoallergic reaction. These photosensitising substances are oral medicines or topical drugs formulated as gels or creams, cosmetics, perfumes, plants or vegetables.

a) Phytophotodermatitis (Sasseville, 2009):

This phototoxic reaction is caused by contact of the skin with photosensitising plants, which may or may not have been ingested.

- Plants that induce phytophotodermatitis (non-exhaustive list):

- Umbelliferae
- Rutaceae
- Moraceae
- Cruciferae
- Ranunculaceae

- Food that induces phytophotodermatitis (non-exhaustive list):

- Fennel, carrots, parsley, dill, celery, angelica
- Lime, lemon
- Figs
- Mustards

b) Berloque dermatitis:

This is induced by contact of the skin with photosensitising cosmetics such as perfumes or certain types of make-up.

c) Drug-induced phototoxicity (Kutlubay et al., 2014; Scheinfeld et al., 2014; Dawe et al., 2014)

Certain drugs sensitise to UV rays following oral or topical administration. This is known as endogenous photosensitisation of the phototoxicity type.

The nails may display pigmentation, or the nail plate may separate from the nail bed, with or without bleeding.

Many drugs may be sensitising, without inducing any actual toxicity, but may, as a result of their effects on the skin, enhance the absorption of UV radiation and thus trigger an increased hypersensitivity to UV radiation. This is looked upon as a case of endogenous photosensitisation of the photoallergy-type, either to UV-A and UV-B radiation.

- Photosensitising drugs inducing phototoxicity-type reactions (non-exhaustive list):

- Oral:

- Antibiotics (cyclins, quinolones)

- Anti-inflammatory drugs (piroxicam, ketoprofen,...)

Anti-epileptic drugs
Psoralens
Etc.
Topical:
Antibiotics (sulfonamides, ...)
Psoralens
Anti-acne medication (tretinoin, benzoyl peroxide ...)
Anti-inflammatory drugs (naproxen, piroxicam...)
Etc.

- Photosensitising drugs inducing photoallergy-type reactions (non-exhaustive list):

Oral:
Anti-tumoral drugs (methotrexate, dacarbazine, ...)
Diuretics (Lasix)
Antianginal drugs (amiodarone, furosemide, ...)
Anti-diabetic drugs
Etc.
Topical:
Antiseptics (chlorhexidine, triclosan,...)
Anti-inflammatory drugs (naproxen, piroxicam...)
Etc.

KEY MESSAGES

The World Health Organization believes that 20 % of cataracts (clouding of the lens), viz. 3 million cases each year, could be due to exposure to UV radiation.

It is strongly advised to regularly wear sunglasses that block UV-A and UV-B radiation.

Photoallergic reactions are abnormal reactions that are triggered by an allergy to one or more types of solar radiation (UV-B and/or UV-A radiation and/or visible light) and take on different shapes.

Some conditions may worsen under the influence of UV radiation.

Photosensitisation reactions can affect anyone from the first exposure to UV radiation, both natural and artificial.

Photosensitising substances are oral medicines or topical drugs formulated as gels or creams, cosmetics, perfumes, plants or vegetables.

V CONCLUSIES EN AANBEVELINGEN

1 Conclusies

De HGR heeft recente gegevens onderzocht om zijn advies over zonnebanken te updaten (HGR 5783, 2000).

Er werd een *ad hoc*-werkgroep opgericht met deskundigen in de volgende disciplines: risicoanalyse, dermatologie, psychologie, oogheelkunde en biofysica. Deze werkgroep heeft de literatuur over de gezondheidseffecten van UV-straling beoordeeld, met speciale aandacht voor zonnebanken voor cosmetische doeleinden.

Uit de literatuur kunnen de volgende conclusies worden getrokken:

- de UV-index is een maat die het effect van de intensiteit van de zonnestraling op de onbeschermdede huid weergeeft. De werkelijke effecten zijn afhankelijk van het huidtype, maar algemeen wordt aanvaard dat bescherming nodig is wanneer de UV-index meer dan 3 bedraagt. In veel landen wordt de huidige en/of verwachte UV-index samen met het weerbericht meegedeeld.
- In vergelijking met andere Europese landen is het gebruik van zonnebanken in België aanzienlijk.
De belangrijkste reden om onder de zonnebank te liggen is om bruin te worden (voor puur cosmetische doeleinden)
Zonnebankgebruikers zijn voornamelijk vrouwelijk en jong (< 35 jaar).
- UV-geïnduceerde huidpigmentatie wordt alleen verkregen via de vorming van pyrimidine-dimeren (d.w.z.: DNA-schade).

Een natuurlijke huidpigmentatie (huidtypes IV tot VI) biedt bescherming tegen huidkanker, terwijl UV-geïnduceerde pigmentatie overeenstemt met een beschermingsfactor (SPF) van niet meer dan 2-4 (vergelijkbaar met zonnecrèmes met een SPF van 2).

UV-geïnduceerde pigmentatie zorgt maar voor een lage fotoprotectie. Blootstelling aan kunstmatige UV-straling voorafgaand aan blootstelling aan de zon moet worden vermeden. Er is dus geen goede reden om zich ter voorkoming van zonnebrand aan kunstmatige UV-straling bloot te stellen.

- Sociologische gegevens bieden een verklaring voor de veranderingen in de gewoonten en gebruiken die verband houden met de verhouding die de mens ten opzichte van de zon aanneemt. Deze verhouding verschilt immers van het ene tijdperk tot het andere en van het ene continent tot het andere en wordt aanzienlijk beïnvloed door de dictaten van de mode. In ons deel van de wereld kwam het bruinen voor het eerst in de jaren 1920 en 1930 in de mode. Deze trend raakte in een stroomversnelling toen het betaald verlof en langere vakanties een realiteit werden. Een bruin uiterlijk werd toen het symbool van een "goede gezondheid". Aangewakkerd door de media leidde deze zoektocht naar zonneshijn logischerwijs tot het ontstaan van zonnebanken. Op een meer psychologisch niveau berust dit verlangen naar een bruin uiterlijk op het zelfbeeld dat men aan anderen wil tonen en dat gekoppeld is aan de "uiterlijke persoon" en het zoeken naar een bepaald ideaal.

Terwijl deze gedrevenheid om aan te trekken en te verleiden typisch haar oorsprong in de narcistische aspecten van onze persoonlijkheid vindt, kan ze echter bij sommige kwetsbaardere individuen ontaarden in een vorm van verslaving. Deze pathologische verslaving aan zonnen, die bekend staat als "tanorexia", is het gevolg van de endorfines die tijdens een chronische blootstelling aan UV-straling vrijkomen. Ze komt hoofdzakelijk voor bij tieners, vooral meisjes, en vaak in combinatie met andere psychiatrische stoornissen zoals angststoornissen, obsessieve-compulsieve stoornissen, stoornis in de lichaamsbeleving en verslavingsziekten.

- Er is momenteel geen bewijs dat een hogere vitamine D-status bij gezonde personen met een normaal dieet in verband kan worden gebracht met gezondheidsvoordelen, d.w.z. het voorkomen van ziekten of het verbeteren van hun uitkomst. Blootstelling aan kunstmatige UV-straling om de vitamine D-status te verhogen is niet efficiënter dan het nemen van orale vitamine D-supplementen, maar gaat wel gepaard met een verhoogd risico op verschillende gezondheidsaandoeningen, waaronder huid- en oogkanker. Het nemen van orale supplementen aan gewone dosissen zoals aanbevolen door de HGR (d.w.z. 10 tot 20 µg per dag) is veilig.
- Uit experimenten met transgene muizen die dragers zijn van één van de belangrijke driver mutaties, blijkt dat UV-bestraling de ontwikkeling van de desbetreffende huidkanker sterk kan bevorderen.

UVA en UVB-straling zijn beide kankerverwekkend.

Recente beoordelingen van biologische gegevens, dierproeven en meta-analyses van de risico's van blootstelling aan UV-straling (inclusief zonnebankgebruik) bevestigen dat er een verband bestaat tussen enerzijds blootstelling aan UV-straling, zowel natuurlijk als onder de zonnebank, en anderzijds het risico op huidkanker, vooral melanoom, en plaveiselcelcarcinoom.

- Volgens ramingen van de effecten van een preventiecampagne tegen huidkanker en die van een algemeen verbod op het gebruik van zonnebanken zouden hierdoor over een periode van 50 jaar bijna 6 000 gevallen van melanoomsterfte kunnen worden vermeden. Bovendien zouden de kosten voor de volksgezondheid dankzij een bewustmakingscampagne met 227,7 miljoen euro kunnen dalen, terwijl een algemeen verbod op zonnebanken in België tot een besparing van € 238 miljoen zou kunnen leiden.
- Heliodermie is de extrinsieke veroudering van de huid die wordt veroorzaakt door intense en/of frequente en cumulatieve blootstelling aan de zon en wordt in verband gebracht met de schadelijke effecten van UV-straling, ook die van zonnebanken. Er bestaan fysiologische mechanismen die het cellulaire DNA herstellen, maar in geval van aanzienlijke herhaalde en cumulatieve dosissen UV-straling zijn deze reparatieprocessen overbelast, waardoor de huid in de loop van de jaren langzaam verandert om naar een precancereuze toestand, of "dysplastische toestand", te evolueren. Actinische keratosen zijn UV-geïnduceerde precancereuze laesies die in ongeveer 5 tot 20 % van de gevallen binnen 10 tot 20 jaar kunnen doorgroeien naar een plaveiselcelcarcinoom.

- De Wereldgezondheidsorganisatie is van oordeel dat 20 % van de gevallen van cataract (lensvertroebeling), nl. 3 miljoen gevallen per jaar, het gevolg zouden kunnen zijn van blootstelling aan UV-straling.
Er wordt sterk aangeraden om regelmatig een zonnebril te dragen die de UVA- en UVB-straling tegenhoudt. Fotoallergische reacties zijn abnormale reacties die worden veroorzaakt door een allergie tegen één of meerdere soorten zonnestraling (UVB en/of UVA-straling en/of zichtbaar licht) en verschillende vormen aannemen.
Bepaalde aandoeningen kunnen onder de invloed van UV-straling verergeren.
Fotosensitisatie kan bij iedereen vanaf de eerste blootstelling aan UV-straling (zowel natuurlijk als kunstmatig) optreden.
Fotosensitiserende stoffen zijn orale geneesmiddelen of topische geneesmiddelen in de vorm van gel of crème, cosmetica, parfums, planten of groenten.
- Bij sommige mensen kan tanorexia, d.w.z. een pathologische verslaving aan het zonnen, ontstaan.

2 Aanbevelingen

Op basis van de solide bewijzen dat UV-straling, met inbegrip van de UV-straling van zonnebanken, huidkanker en andere huid- en oogziekten veroorzaakt, en rekening houdend met het feit dat

- er voor het opwekken van huidkanker geen drempelniveau bestaat voor UV-straling en dus ook geen veilige blootstellingsgrens;
- de risico's van blootstelling aan UV-straling zwaarder wegen dan de voordelen, zoals de aanmaak van vitamine D;
- een minimale blootstelling aan natuurlijke UV-straling doorgaans voldoende is om de nodige hoeveelheden vitamine D aan te maken;
- het gebruik van zonnebanken geen goede oplossing is om een vitamine D-tekort te verhelpen;
- zonnebanken permanent beschikbaar zijn, wat ertoe aanzet om deze herhaaldelijk en overmatig te gebruiken.
De blootstelling aan natuurlijke UV-straling kan aan de hand van beschermende maatregelen worden beperkt, maar niet volledig worden uitgesloten.

De HGR adviseert om zonnebanken en alle bronnen van kunstmatige UV-straling die beschikbaar zijn voor het publiek te verbieden, met als doel het risico op huidkanker te verminderen.

Tijdens de overgangperiode tot het verbod op zonnebanken effectief is, en ook achteraf, beveelt de HGR aan om de algemene bevolking op een objectieve en transparante manier in te lichten over de risico's van UV-straling in het algemeen (en ook over de risico's van de UV-straling die door zonnebanken wordt geproduceerd), en over de redenen achter het verbod op zonnebanken. De bedoeling is om zo een verschuiving naar een verhoogde blootstelling aan natuurlijke UV-straling te vermijden. Bovendien is het ook belangrijk om in deze voorlichtingscampagne objectieve informatie op te nemen over de positieve effecten van incidentele blootstelling aan de zon, bv. de aanmaak van vitamine D.

3 Recommendations

Based on the well-established evidence that UVR, including UVR emitted by sunbeds, induces skin cancer and other cutaneous and ocular diseases, and taking into account that:

- There is no threshold level of UVR for the induction of skin cancer and hence no safe limit for exposure to UV radiation;
- The risks of UVR exposure outweigh the benefits such as the induction of vitamin D;
- Minimal exposure to natural UVR is in general sufficient to induce sufficient amounts of vitamin D;
- Sunbed use is not the way to replenish vitamin D deficiencies;
- Sunbed use is continuously available and invites for repetitive and excessive use. Exposure to natural UVR can be limited by protective measures but cannot be excluded completely.

The SHC recommends to ban the sunbeds and all artificial UVR devices that are available to the public, with the aim of reducing the risk for skin cancer.

In the interim period to the effective ban of sunbeds, and also afterwards, in order to avoid a shift towards an increased exposure to natural UVR, the SHC further recommends to inform the general public in an objective and transparent way about the risks linked to UVR in general (and including the risks of UVR emitted by sunbeds), and about the rationale to ban the sunbeds. Moreover, it is also important to include objective information about the beneficial effects of casual exposure to the sun e.g. vitamin D production, in this information campaign.

VI REFERENCES

- ACGIH. Ultraviolet Radiation: TLV Physical Agents 7th Edition Documentation. Cincinnati, OH: ACGIH; 2010.
- Agar NS, Halliday GM, Barnetson RS, Ananthaswamy HN, Wheeler M, & Jones AM. The basal layer in human squamous tumors harbors more uva than uvb fingerprint mutations: A role for uva in human skin carcinogenesis. *Proc Natl Acad Sci USA*, 2004; 101(14), 4954-4959.
- Alberg AJ. "A melanoma epidemic in iceland: Possible influence of sunbed use". *Am J Epidemiol* 2011; 73(7), 845.
- Albert MR, Ostheimer KG. The evolution of current medical and popular attitudes toward ultraviolet light exposure: part 1. *J Am Acad Dermatol* 2002; 47(6):930-7.
- Albert MR, Ostheimer KG. The evolution of current medical and popular attitudes toward ultraviolet light exposure: part 2. *J Am Acad Dermatol* 2003a; 48(6):909-18.
- Albert MR, Ostheimer KG. The evolution of current medical and popular attitudes toward ultraviolet light exposure: part 3. *J Am Acad Dermatol* 2003b; 49(6):1096-106.
- Amrein K, Schnedl C, Holl A, Riedl R, Christopher KB, Pachler C et al. Effect of high-dose vitamin d3 on hospital length of stay in critically ill patients with vitamin d deficiency: The vitdal-icu randomized clinical trial. *JAMA* 2014; 312(15), 1520-1530
- Ashrafioun & Bonar EE. Tanning addiction and psychopathology: Further evaluation of anxiety disorders and substance abuse. *J Am Acad Dermatol*. 2014; 70(3):473-80.
- Autier P & Boyle P. Artificial ultraviolet sources and skin cancers: Rationale for restricting access to sunbed use before 18 years of age. *Nat Clin Pract Oncol* 2008; 5(4):178-179.
- Autier P & Dore JF. Influence of sun exposures during childhood and during adulthood on melanoma risk. Epimel and eortc melanoma cooperative group. European organisation for research and treatment of cancer. *Int J Cancer* 1998; 77(4): 533-537.
- Autier P, Boniol M, Pizot C, & Mullie P. Vitamin d status and ill health - author's reply. *Lancet Diabetes Endocrinol* 2014a; 2(4):275-276.
- Autier P. Epidemiological evidence that uva is involved in the genesis of melanoma Sunbed use, sunscreen use, childhood sun exposure, and cutaneous melanoma. 2011
- Autier P. Answer to the commentary: A sunbed epidemic? *Am J Epidemiol*. 2010; 172 (7): 771-772.
- Belgian Cancer foundation. Kankerbarometer – Resultaten 2015. Ipsos Public Affairs. The Social Research and Corporate Reputation Specialists. Ipsos, 2015b. (http://www.kanker.be/sites/default/files/KankerbarometerSTK_resultaten_2015.pdf).
- Belgian Cancer foundation. Tracking survey 'knowledge, attitude and behavior of the Belgians towards UV'. Ipsos Public Affairs. The Social Research and Corporate Reputation Specialists. Ipsos, 2015a.

Berking C, Takemoto R, Binder RL, et al. Photocarcinogenesis in human adult skin grafts. *Carcinogenesis* 2002; 23(1):181-7.

Bolland MJ, & Grey A. Are trials of vitamin d with mortality as an endpoint really needed? *BMJ* 2014a; 349:g4452.

Bolland MJ, Grey A, Gamble GD, & Reid IR. Are more trials of vitamin d supplementation needed for skeletal, vascular or cancer outcomes? A trial sequential meta-analysis. *Lancet Diabetes Endocrinol* 2014b.

Bolland MJ, Grey A, Gamble GD, & Reid IR. (2014c). The effect of vitamin d supplementation on skeletal, vascular, or cancer outcomes: A trial sequential meta-analysis. *The Lancet Diabetes & Endocrinol* 2014c; 2(4):307-320.

Boniol M, Autier P, Boyle P, Gandini S. Cutaneous melanoma attributable to sunbed use: systematic review and meta-analysis. *BMJ* 2012; 345:e4757.

Brenner M, Hearing VJ. The protective role of melanin against UV damage in human skin. *Photochem Photobiol* 2008; 84(3):539-49.

Bruggers JHA, de Jong WE, Bosnjakovic BFM, Passchier WF. Use of artificial tanning equipment in the Netherlands. In: Passchier WF et al., editors. *Human exposure to ultraviolet radiation Risk and regulations Excerpta Medica International Congress Series 744*. Amsterdam: Elsevier Science Publishers 1987; p. 235-9.

Calderone DC, Fenske NA. The clinical spectrum of actinic elastosis. *J Am Acad Dermatol*. 1995; 32:1016–1024

Carter S. *Rise and Shine: Sunlight, Technology and Health*. Oxford: Berg Publishers; 2007. ISBN-13: 978-1845201319.

Ceburkov O, Gollnick H. Photodynamic therapy in dermatology. *Eur J Dermatol* 2000; 10(7):568-75.

Choudhary S, Nouri K, Elsaie ML. Photodynamic therapy in dermatology: a review. *Lasers Med Sci*. 2009; 24(6):971-80.

Chowdhury R, Kunutsor S, Vitezova A, Oliver-Williams C, Chowdhury S, Kieft-de-Jong JC et al. Vitamin D and risk of cause specific death: systematic review and meta-analysis of observational cohort and randomised intervention studies. *BMJ*. 2014; 348:g1903.

Commission Internationale de l'Éclairage. *International Lighting Vocabulary*. Vienna: CIE Central Bureau; 2009. Publication CIE DS 017.2:2009.

Commission Internationale de l'Éclairage. *Recommendations on minimum levels of solar UV exposure*. Vienna: Commission Internationale de l'éclairage; 2011. Technical Report CIE 201:2011.

Danish Cancer Society and TrygFonden. *Sun survey 2012*. Available online (http://www.cancer.dk/dyn/resources/File/file/7/137/1385246568/sunsurvey_2012.pdf)

Davie M, & Lawson DE. Assessment of plasma 25-hydroxyvitamin d response to ultraviolet irradiation over a controlled area in young and elderly subjects. *Clin Sci (Lond)* 1980; 58(3):235-242.

Dawe RS, Ibbotson SH. Drug-induced photosensitivity. *Dermatol Clin*. 2014 Jul; 32(3):363-8.

De Mol J. Le dommage esthétique in "Le dommage psychique: du traumatisme à l'expertise"; Larcier, 2012, 302 p.

Diffey BL, Roscoe AH. Exposure to solar ultraviolet radiation in flight. *Aviat Space Environ Med* 1990; 61(11):1032-5.

Diffey BL. Observations on the use of UV-A sunbeds for cosmetic tanning. In: Passchier WF et al., editors. Human exposure to ultraviolet radiation Risk and regulations Excerpta Medica International Congress Series 744. Amsterdam: Elsevier Science Publishers; 1987. p. 241-6.

Diffey BL. Sources and measurement of ultraviolet radiation. *Methods* 2002; 28(1):4-13.

Diffey BL. The risk of squamous cell carcinoma in women from exposure to UVA lamps used in cosmetic nail treatment. *Br J Dermatol* 2012; 167(5):1175-8.

El Ghissassi F, Baan R, Straif K, Grosse Y, Secretan B, Bouvard V et al. A review of human carcinogens—Part D: radiation. *Lancet Oncol* 2009; 10:751-2.

Erdmann F, Lortet-Tieulent J, Schüz J, Zeeb H, Greinert R, Breitbart EW et al. International trends in the incidence of malignant melanoma 1953-2008—are recent generations at higher or lower risk? *International Journal of Cancer* 2013; 132(2): 385-400.

Fell GL, Robinson KC, Mao J, Woolf CJ, Fisher DE. Skin β -Endorphin Mediates Addiction to UV Light. *Cell* 2014; 157(7):1527-34.

Fitzpatrick's Dermatology in General Medicine, 5th ed

Gerber B, Mathys P, Moser M, Bressoud D, & Braun-Fahrlander C. Ultraviolet emission spectra of sunbeds. *Photochem Photobiol* 2002; 76(6): 664-668.

Gezondheidsraad. UV straling en zonnebanken. Den Haag: Gezondheidsraad; 2009. Publicatienr. 2009/11. Internet: <http://www.gezondheidsraad.nl/nl/adviezen/gezonde-leefomgeving/briefadvies-uv-straling-en-zonnebanken>

Gezondheidsraad: Commissie UV straling. UV straling: blootstelling van de mens aan ultraviolette straling [Ultraviolet Radiation human exposure to ultraviolet radiation]. Den Haag: Gezondheidsraad; 1986. Publicatie nr. 1986/09. Internet: <http://www.gezondheidsraad.nl/nl/taak-werkwijze/werkterrein/gezonde-leefomgeving/uv-straling-blootstelling-van-de-mens-aan>,

Goetze S, Elsner P. Solar urticaria *J Dtsch Dermatol Ges*. 2015; 13(12):1250-3.

Green AC. Epidemiology of actinic keratoses. *Curr Probl Dermatol* 2015; 46:1-7.

Gruber-Wackernagel A, Byrne SN, Wolf P. Polymorphous light eruption: clinic aspects and pathogenesis, *Dermatol Clin* 2014; 32(3):315-34.

Harrington CR, Beswick TC, Leitenberger J, Minhajuddin A, Jacobe HT, Adinoff B. Addictive-like behaviours to ultraviolet light among frequent indoor tanners. *Clin Exp Dermatol* 2011; 36(1):33-8.

Hart P, Norval M, Reeve V. Introduction to the themed issue 'The health benefits of UV radiation exposure through vitamin D production or non-vitamin D pathways'. *Photochem Photobiol Sci* 2017; 16(3): 281-282.

Harvey NC, & Cooper C. Vitamin d: Some perspective please. *BMJ* 2012; 345:e4695.

Heckman CJ, Darlow S, Kloss JD, Cohen-Filipic J, Manne SL, Munshi et al. Measurement of tanning dependence. *J Eur Acad Dermatol Venereol* 2014; 28(9):1179-85.

Hess AF, Unger LF. Cure of infantile rickets by sunlight. *J Am Med Ass* 1921; 77:39.

HGR – Hoge Gezondheidsraad. Advies inzake zonnebanken en UV straling. Advies no 5783, December 2000.

HGR - Hoge Gezondheidsraad. UV nagellampen. Brussel: HGR; 2013. Advies no 9102.

HGR – Hoge Gezondheidsraad. Recommendations nutritionnelles pour la Belgique 2016. Advies no 9285.

Hillhouse JJ, Baker MK, Turrisi R, Shields A, Stapleton J, Jain S et al. Evaluating a measure of tanning abuse and dependence. *Arch Dermatol* 2012; 148(7):815-9.

Holly EA, Aston DA, Char DH, Kristiansen JJ, & Ahn DK. Uveal melanoma in relation to ultraviolet light exposure and host factors. *Cancer Res* 1990; 50(18): 5773-5777.

IARC working group. (2006). Exposure to artificial uv radiation and skin cancer. In World Health Organization International Agency for Research on Cancer (Ed.). Lyon, France.

IARC. (1992). IARC monographs on the evaluation of carcinogenic risks to humans. Solar and ultraviolet radiation IARC Monogr Eval Carcinog Risks Hum (1992/01/01 ed., Vol. 55, pp. 1-316). Lyon, France: International Agency for Research on Cancer.

International Commission on Non-Ionizing Radiation Protection. ICNIRP Statement - Protection of Workers against Ultraviolet Radiation. *Health Phys* 2010; 99(1):66-87.

Kourosh AS, Harrington CR, Adinoff B. Tanning as a behavioral addiction. *Am J Drug Alcohol Abuse* 2010; 36(5):284-90.

Kuhn A, Beissert S. Photosensitivity in lupus erythematosus. *Autoimmunity*. 2005; 38(7):519-29.

Kutlubay Z, Sevim A, Engin B, Tüzün Y. Photodermatoses, including phototoxic and photoallergic reactions (internal and external) *Clin Dermatol*. 2014; 32(1):73-9.

Lagunova Z, Porojnicu AC, Aksnes L, Holick MF, Iani V, Bruland OS et al. Effect of vitamin d supplementation and ultraviolet b exposure on serum 25-hydroxyvitamin d concentrations in healthy volunteers: A randomized, crossover clinical trial. *Br J Dermatol* 2013; 169(2):434-440.

Langdahl JH, Schierbeck LL, Bang UC, Jensen JE. Changes in serum 25-hydroxyvitamin D and cholecalciferol after one whole-body exposure in a commercial tanning bed: a randomized study. *Endocrine*. 2012; 42(2):430-5.

Lava SA, Simonetti GD, Ragazzi M, Guarino Gubler S, Bianchetti MG. Juvenile spring eruption: an outbreak report and systematic review of the literature. *Br J Dermatol*. 2013; 168(5):1066-72.

Manson JE, Bassuk SS. Vitamin D research and clinical practice: at a crossroads. *JAMA* 2015; 313(13):1311-2.

Meyer HE, Holvik K, & Lips P. Should vitamin d supplements be recommended to prevent chronic diseases? *BMJ (Online)*, 2015; 350.

Miller SA, Hamilton SL, Wester UG, & Cyr WH. An analysis of uva emissions from sunlamps and the potential importance for melanoma. *Photochem Photobiol* 1998; 68(1):63-70.

Miyamura Y, Coelho SG, Schlenz K, Batzer J, Smuda C, Choi W, Brenner et al. The deceptive nature of UVA tanning versus the modest protective effects of UVB tanning on human skin. *Pigment Cell Melanoma Res.* 2011; 24(1):136-47.

Mosher CE, Danoff-Burg S. Addiction to Indoor Tanning. Relation to Anxiety, Depression and Substance Use. *Archives of Dermatology*, 2010; 146(4):412-7.

Mouret S, Baudouin C, Charveron M, Favier A, Cadet J and Douki T. Cyclobutane pyrimidine dimers are predominant DNA lesions in whole human skin exposed to uva radiation. *Proc Natl Acad Sci USA* 2006; 103(37):13765-13770.

Nakamura M, Morita A, Seit  S, Haarmann-Stemmann T, Grether-Beck S, Krutmann J. Environment-induced lentigines: formation of solar lentigines beyond ultraviolet radiation. *Exp Dermatol.* 2015; 24(6):407-11.

Nilsen LT, Aalerud TN, Hannevik M, Veierod MB. High UV-A exposure from sunbeds. *Pigment Cell Melanoma Res* 2012; 25(5):639-40.

Nilsen LT, Aalerud TN, Hannevik M, Veierod MB. UVB and UVA irradiances from indoor tanning devices. *Photochem Photobiol Sci* 2011; 10(7):1129-36.

Noonan FP, Recio JA, Takayama H, Duray P, Anver MR, Rush WL et al. Neonatal sunburn and melanoma in mice. *Nature* 2001; 413(6853):271-272.

Noonan FP, Recio JA, Takayama H, et al. Neonatal sunburn and melanoma in mice. *Nature* 2001; 413(6853):271-2.

Noonan FP, Zaidi MR, Wolnicka-Glubisz A, Anver MR, Bahn J, Wielgus A et al. Melanoma induction by ultraviolet a but not ultraviolet b radiation requires melanin pigment. *Nat Commun* 2012; 3:884.

Onoue S, Seto Y, Sato H, Nishida H, Hirota M, Ashikaga T, Api AM, et al. Chemical photoallergy: photobiochemical mechanisms, classification, and risk assessments, *J Dermatol Sci.* 2017 ; 85(1):4-11.

Ontario Government. Ultraviolet Radiation in the Workplace. Toronto: Ontario Ministry of Labour; 2009 March. Internet: <http://www.labour.gov.on.ca/english/hs/pubs/uvradiation/index.php>, accessed 19-01-2016.

Ortonne JP. The effects of ultraviolet exposure on skin melanin pigmentation. *J Int Med Res.* 1990; 18 Suppl 3:8C-17C.

Palma-Carlos AG, Palma-Carlos ML. Solar urticaria and porphyriasa . *Eur Ann Allergy Clin Immunol.* 2005; 37(1):17-20.

Petit A, Karila L, Lejoyeux M. Quel cadre nosographique pour le bronzage excessif ? *Encéphale* 2014; 40(2):174-9.

Petit A, Lejoyeux M. Peut-on parler d'addiction pour le bronzage excessif en cabine? *Rev Med Liège* 2013; 68(5-6):315-20.

Petit A, Richoux C, Lejoyeux M. L'excès de bronzage constitue-t-il une nouvelle forme de dépendance. *Alcoologie et Addictologie*, 2011, 33(3):259-64.

Philips. Meulemans CCE, editor. Modern suntanning methods. Guidelines for professional users. Eindhoven, The Netherlands: Nederlands Philips Bedrijven, Lighting Division; 1987 January. No 1/87.

Pil L, Hoorens I, Vossaert K, Kruse V, Tromme I, Speybroeck N et al., Burden of skin cancer in Belgium and cost-effectiveness of primary prevention by reducing ultraviolet exposure. *Prev Med.* 2016; 93:177-182.

Prosafe. Joint Market surveillance report, Joint action best practices, Prosafe, 2012.

Radiation. IARC Monogr Eval Carcinog Risks Hum 2012; 7:303.

Rodríguez-Carreón AA, Rodríguez-Lobato E, Rodríguez-Gutiérrez G, Cuevas-González JC, Mancheno-Valencia A, et al. Actinic Prurigo. *Skinmed.* 2015; 13(4):287-95.

Sallander E, Wester U, Bengtsson E, & Wiegleb Edström D. Vitamin d levels after uvb radiation: Effects by uva additions in a randomized controlled trial. *Photodermatology, Photoimmunology & Photomedicine* 2013; 29(6):323-329.

Sasseville D. Clinical patterns of phytodermatitis. *Dermatol Clin* 2009; 27(3):299-308.

SCHEER, Opinion on biological effects of ultraviolet radiation relevant to health with particular reference to sunbeds for cosmetic purposes, 17 november 2016

Scheinfeld NS, Chernoff K, Derek Ho MK, Liu YC. Drug-induced photoallergic and phototoxic reactions - an update *Expert Opin Drug Saf.* 2014; 13(3):321-40.

Schneider S, Diehl K, Bock C, Schlüter M, Breibart AW, Volkmer B et al. Sunbed use, user characteristics, and motivations for tanning: results from the German population-based SUN-Study 2012. *JAMA Dermatol* 2013; 149(1):43-49.

Seddon JM, Gragoudas ES, Glynn RJ, Egan KM, Albert DM, & Blitzer PH. Host factors, uv radiation, and risk of uveal melanoma. A case-control study. *Arch Ophthalmol* 1990; 108(9): 1274-1280.

The Skin Cancer Foundation's Position Regarding UVR-Emitting Nail Lamps New York: The Skin Cancer Foundation; 2013 March 17. Internet: <http://www.skincancer.org/media-and-press/press-release-2013/nail-lamps>, accessed 01-09-2014.

Theodoratou E, Tzoulaki I, Zgaga L, & Ioannidis JPA. Vitamin d and multiple health outcomes: Umbrella review of systematic reviews and meta-analyses of observational studies and randomised trials. *BMJ* 2014; 348.

Thieden E, Jørgensen HL, Jørgensen NR, Philipsen PA, Wulf HC. Sunbed radiation provokes cutaneous vitamin D synthesis in humans--a randomized controlled trial. *Photochem Photobiol*. 2008; 84(6):1487-92.

Tierney P, de Grujil FR, Ibbotson S, & Moseley H. Predicted increased risk of squamous cell carcinoma induction associated with sunbed exposure habits. *Br J Dermatol* 2015; 173(1): 201-208.

Tierney P, Ferguson J, Ibbotson S, Dawe R, Eadie E, Moseley H. Nine out of 10 sunbeds in England emit ultraviolet radiation levels that exceed current safety limits. *Br J Dermatol* 2013; 168(3):602-8.

Trakatelli M, Barkitzi K, Apap C, Majewski S, De Vries E. Skin cancer risk in outdoor workers: a European multicenter case-control study. *J Eur Acad Dermatol Venereol*. 2016; 30 Suppl 3:5-11.

Tran TT, Schulman J, Fisher DE. UV and pigmentation: molecular mechanisms and social controversies. *Pigment Cell Melanoma Res*. 2008; 21(5):509-16.

Tucker MA, Hartge P, & Shields JA. Epidemiology of intraocular melanoma. *Recent Results Cancer Res* 1986; 102:159-165.

Uv radiation & skin cancer. The science behind age restrictions for tanning beds. *Environmental Health Perspective* 2012; 120(8):A308-313.

Vajdic CM, Krickler A, Giblin M, McKenzie J, Aitken J, Giles GG, Armstrong BK. Sun exposure predicts risk of ocular melanoma in australia. *Int J Cancer* 2002; 101(2):175-182.

Warthan MM, Uchida T, Wagner RF. UV light tanning as a type of substance-related disorder. *Arch Dermatol* 2005; 141(8):963-6.

Wehner MR, Shive ML, Chren MM, Han J, Qureshi AA, Linos E. Indoor tanning and non-melanoma skin cancer: Systematic review and meta-analysis. *BMJ* 2012; 345:e5909.

Werkgroep 'Relatie kanker, zonlicht en vitamine D' van de Signaleringscommissie Kanker van KWF Kankerbestrijding. De relatie tussen kanker, zonnestraling en vitamine D. Amsterdam: KWF Kankerbestrijding; 2010 augustus. Internet: <http://www.kwf.nl/SiteCollectionDocuments/Rapport-De-relatie-tussen-kanker-zonnestraling-en-vitamineD.pdf>, accessed 05-06-2014.

Whiteman DC, Whiteman CA, & Green AC. Childhood sun exposure as a risk factor for melanoma: A systematic review of epidemiologic studies. *Cancer Causes Control* 2001; 12(1), 69-82.

Wolnicka-Glubisz A, Noonan FP. Neonatal susceptibility to UV induced cutaneous malignant melanoma in a mouse model. *Photochem Photobiol Sci* 2006; 5(2):254-60.

VII COMPOSITION OF THE WORKING GROUP

The composition of the Committee and that of the Board as well as the list of experts appointed by Royal Decree are available on the following website: [composition and mode of operation](#).

All experts joined the working group *in a private capacity*. Their general declarations of interests as well as those of the members of the Committee and the Board can be viewed on the SHC website (site: [conflicts of interest](#)).

The following experts were involved in drawing up and endorsing this advisory report. The working group was chaired by **Olivier VANHOOTEGHEM**; the scientific secretary was Veerle MERTENS.

ADANG Dirk	Health and environment	UCL
AUTIER Philippe	Skin cancer, epidemiology	IPRI
BOONEN Brigitte	Skin cancer prevention	Belgian Cancer Foundation
BROCHEZ Lieve	Skin cancer	UGent
DE BACKER Hugo	Weather instruments	Royal Meteorological Institute of Belgium
DE MOL Jacques	Psychology	
DEL MARMOL Véronique	Skin cancer, epidemiology	ULB
KOPPEN Carina	Ophthalmology, visual optics and visual revalidation	UAntwerpen
VANHOOTEGHEM Olivier	Surgical dermatology	CHU UCL Namur

The following experts were heard:

VAN RONGEN Eric	Radiation expert	Health Council Netherlands
DE GRUIJL Frank	Photobiology/physics	Leiden University
PASSCHIER Wim	Health risk analysis	Maastricht University

The following administrations and/or ministerial cabinets were heard:

MEUNIER Joëlle	Head of the cell Cosmetics	SPF SPSCAE
JACQUES Sarah	Regulation expert	FPS Economy
PASTEELS Karine	Technical expert	SPF SPSC
VERCKENS Bram	Regulation expert	FPS Economy

The following firms/associations/etc. were heard:

BOECKX John	Chair	BESKO
DELGOFFE Daniel	Technical advisor	UNEB
GYS-BEHETS Francine	Honorary chair	BESKO
SALEMBIER Nadine	Chair	UNEB

The standing working group Cosmetics has endorsed the advisory report. The standing working group was chaired by **Albert DE MEY**; the scientific secretary was Veerle MERTENS.

The standing working group of Non-Ionising Radiation has endorsed the advisory report. The standing working group was chaired by **Luc VERSCHAEVE**; the scientific secretary was Eric JADOUL.

VIII APPENDIXES

1 Appendix 1: Definition and interpretation of the UV index

Figure a explains how the UV-index is calculated. The yellow line shows the intensity of the solar radiation on a horizontal surface on a typical summer day in Belgium as a function of wavelengths in the UV-B and UV-A range (mind the logarithmic scale). The blue line shows a typical spectrum reaching the surface at a mid-latitudinal location at noon during summer. The cyan line shows the action spectrum for erythema in human skin (<http://www.cie.co.at/publ/abst/s007.html>). Multiplying this with the measured spectrum gives the red line. The UV-index is obtained by integrating this curve between 280 and 400 nm and multiplying it by 40.




Figure a: Definition of the UV-index

The meaning of the index (for typical European skin) is explained in table a.

Table a Relationship between UV-index, UV-intensity and (European) skin burning¹⁰

UV-Index	UV-Intensity	Skin burns
0-2	low	almost not
2-4	moderate	slowly
4-6	moderately high	easily
6-8	high	fast
8-10	very high	very fast
a>10	extreme	almost immediately

This index can also be forecasted¹¹ for clear sky conditions using radiative transfer models. Measurements of the UV index are also available^{12 and 13}.

In Belgium the UV index reaches values of 7-8 in early summer at solar noon (corresponding to approximately 13:45 local time). On the Canary Islands, for example, the maximum value can go up to 12 and even higher. In high-mountain regions (e.g. Himalaya in Nepal), where UV radiation reflected from the snow, values of 20 or higher are sometimes observed. As mentioned in section 1.2, sunbeds may expose users to effective erythemal intensities of maximum 0.3 W/m². Following the WMO/WHO definition, this corresponds to a UV-index of 12.

¹⁰Classification according to the Royal Meteorological Institute of Belgium (adapted from WHO, 2002).

¹¹These forecasts are distributed together with the weather forecasts (<http://www.meteo.be/meteo/view/nl/123504-Uv-index.html>)

¹²<http://www.meteo.be/meteo/view/nl/123504-Uv-index.htm>

2 Appendix 2: Results of Euromelanoma survey (31 countries)


2.1.1 EU population

2.1.2 Sunbed use

	Valid N	N	%
Period 2009-2014	218,433	19,467	8.9%
2015	44,212	4,416	10.0%


	Screenees	Percentage
No	33,775	87,8%
Yes	3,303	8,6%
Unknown	1,406	3,7%
Total	38,484	100%

	Mean	Standard Deviation	Median	Percentile 25	Percentile 75
Belgium	8.35	6.79	6.00	3.00	10.00
Bosnia H	3.32	2.62	3.00	1.00	4.00
Croatia	4.97	3.67	4.00	3.00	5.50
Denmark	10.23	9.21	9.00	3.00	17.50
Estonia	4.33	3.30	3.00	2.00	5.00
Fyro Macedonia	3.88	4.51	3.00	1.50	4.00
Georgia					
Germany	9.21	6.69	9.50	4.00	11.00
Greece	2.85	2.40	2.00	1.00	3.00
Hungary	7.23	4.95	6.00	3.00	10.00
Ireland	9.20	9.88	3.00	2.00	20.00
Latvia	6.18	4.66	5.00	3.00	8.00
Lithuania	4.15	3.44	3.00	2.00	5.00
Malta	1.00	.	1.00	1.00	1.00
Poland	7.45	5.06	6.00	3.00	10.00
Portugal	4.65	4.58	3.50	2.00	5.50
Romania	2.84	2.49	2.00	1.00	3.00
Russia	4.07	3.32	3.00	2.00	5.00
Serbia	3.23	2.66	2.00	2.00	4.00
Sweden	9.11	7.93	7.00	3.00	11.00
Ukraine	3.04	2.21	2.00	1.00	4.00



> 15%: Belgium, Hungary, Latvia, Lithuania, Poland

2.1.3 Sunbed use (females < 35y)



> 20%: Belgium, Croatia, Denmark, Fyromacedonia, Estonia, Germany, Hungary, Latvia, Lithuania, Poland

< 10%: Georgia, Greece, Malta, Portugal, Romania

2.2 Belgian population

2.2.1 Sunbed use (2015)

	Valid N	N	%	< 20 sessions	≥ 21 sessions
15 - 65 years of age	1427	233	16.3%	93.6%	6.4%
18 – 35 years of age	404	66	16.3%	92.4%	7.6%

	Women	Men				
	%	< 20 sessions	≥ 21 sessions	%	< 20 sessions	≥ 21 sessions
15 - 65 years of age	21.7%	20.4%	1.3%	8.7%	8.0%	0.7%
18 – 35 years of age	20.8%	19.7%	1.1%	6.4%	4.8%	1.6%

2.2.2 Number of years of use (2015)

	Mean	Standard Deviation	Median	Percentile 25	Percentile 75
15 - 65 years of age	7.4	7.2	5	2	10
18 – 35 years of age	5.4	4.1	5	2	10

2.2.3 Evolution of sunbed use (2009 – 2015)

		Valid N	N	%	< 20 sessions	≥ 21 sessions
Total	Period 2009-2014	9,901	2,207	22.3%	90.0%	10.0%
	2009	2,605	658	25.3%	89.4%	10.6%
	2010	1,131	237	21.0%	86.1%	13.9%
	2011	1,711	389	22.7%	89.7%	10.3%
	2012	1,652	388	23.5%	91.2%	8.8%
	2013	1,136	217	19.1%	89.4%	10.6%
	2014	1,666	318	19.1%	93.7%	6.3%
15 - 65 years of age	2015	1,723	248	14.4%	93.5%	6.5%
	Period 2009-2014	8,763	2,102	24.0%	90.0%	10.0%
	2009	2,181	619	28.4%	89.0%	11.0%
	2010	1,012	232	22.9%	85.8%	14.2%
	2011	1,592	377	23.7%	89.7%	10.3%

	2012	1,471	372	25.3%	91.1%	8.9%
	2013	1,036	203	19.6%	90.1%	9.9%
	2014	1,471	299	20.3%	94.0%	6.0%
	2015	1,427	233	16.3%	93.6%	6.4%
18 – 35 years of age	Period 2009-2014	2,888	775	26.8%	89.0%	11.0%
	2009	632	202	32.0%	89.1%	10.9%
	2010	397	115	29.0%	82.6%	17.4%
	2011	624	161	25.8%	87.0%	13.0%
	2012	440	122	27.7%	88.5%	11.5%
	2013	302	61	20.2%	91.8%	8.2%
	2014	493	114	23.1%	97.4%	2.6%
	2015	404	66	16.3%	92.4%	7.6%

2.2.4 Evolution of number of years of use

	Mean	Standard Deviation	Median	Percentile 25	Percentile 75
Period 2009-2014	7.5	6.6	5	3	10
2009	6.6	6.4	5	2	10
2010	7.4	6.8	5	3	10
2011	7.9	6.1	5	3	10
2012	8.4	6.8	6	3	10
2013	8.4	7.2	6	3	10
2014	7.8	6.9	5	3	10
2015	7.3	7.1	5	2	10

Over de Hoge Gezondheidsraad (HGR)

De Hoge Gezondheidsraad is een federaal adviesorgaan waarvan de FOD Volksgezondheid, Veiligheid van de Voedselketen en Leefmilieu het secretariaat verzekert. Hij werd opgericht in 1849 en geeft wetenschappelijke adviezen i.v.m. de volksgezondheid aan de ministers van Volksgezondheid en van Leefmilieu, aan hun administraties en aan enkele agentschappen. Hij doet dit op vraag of op eigen initiatief. De HGR probeert het beleid inzake volksgezondheid de weg te wijzen op basis van de recentste wetenschappelijke kennis.

Naast een intern secretariaat van een 25-tal medewerkers, doet de Raad beroep op een uitgebreid netwerk van meer dan 500 experts (universiteitsprofessoren, medewerkers van wetenschappelijke instellingen, praktijkbeoefenaars, enz.), waarvan er 300 tot expert van de Raad zijn benoemd bij KB; de experts komen in multidisciplinaire werkgroepen samen om de adviezen uit te werken.

Als officieel orgaan vindt de Hoge Gezondheidsraad het van fundamenteel belang de neutraliteit en onpartijdigheid te garanderen van de wetenschappelijke adviezen die hij aflevert. Daartoe heeft hij zich voorzien van een structuur, regels en procedures die toelaten doeltreffend tegemoet te komen aan deze behoeften bij iedere stap van het tot stand komen van de adviezen. De sleutelmomenten hierin zijn de voorafgaande analyse van de aanvraag, de aanduiding van de deskundigen voor de werkgroepen, het instellen van een systeem van beheer van mogelijke belangenconflicten (gebaseerd op belangenverklaringen, onderzoek van mogelijke belangenconflicten en een Commissie voor Deontologie) en de uiteindelijke validatie van de adviezen door het College (eindbeslissingsorgaan van de HGR, samengesteld uit 40 leden van de pool van benoemde experts). Dit coherent geheel moet toelaten adviezen af te leveren die gesteund zijn op de hoogst mogelijke beschikbare wetenschappelijke expertise binnen de grootst mogelijke onpartijdigheid.

Na validatie door het College worden de adviezen overgemaakt aan de aanvrager en aan de minister van Volksgezondheid en worden ze gepubliceerd op de website (www.hgr-css.be). Daarnaast wordt een aantal onder hen gecommuniceerd naar de pers en naar bepaalde doelgroepen (beroepsbeoefenaars in de gezondheidssector, universiteiten, politiek, consumentenorganisaties, enz.).

Indien u op de hoogte wilt blijven van de activiteiten en publicaties van de HGR kunt u een mail sturen naar info.hgr-css@health.belgium.be.